

King County Medical Examiner's Office Annual Report 2013



Public Health
Seattle & King County





King County Medical Examiner's Office
 Harborview Medical Center
 325 Ninth Ave, Box 359792
 Seattle, Washington 98104

Phone: 206-731-3232 | Fax: 206-731-8555
 TTY Relay: 711
www.kingcounty.gov/health/examiner

Patty Hayes
PATTY HAYES

Interim Director of Public Health
 Public Health – Seattle & King County

Richard Harruff
RICHARD HARRUFF, MD, PhD

Chief Medical Examiner
 King County Medical Examiner's Office
 Public Health – Seattle & King County



Public Health
 Seattle & King County



2013 Annual Report

DEDICATION

We recognize that each case in this report represents the death of a person whose absence is grieved by friends and relatives. These deaths also represent a loss to our community. As those responsible for investigating these deaths, we dedicate this report to the memory of those lost and to those who have suffered the loss of a friend or relative.

Table of Contents

Foreword	2	Manner of death: Undetermined	66
Executive summary.....	3	Traffic deaths	72
Description and purpose	5	Deaths due to drugs and poisons	81
Mission statement	7	Listed drug names	84
Explanation of data.....	8	Deaths due to firearms	95
Medical Examiner cases in 2013.....	10	Causes of deaths in children and youth.....	99
Manner of death in 2013	15	Organ donation	103
Out of County cases in 2013	22	Disposition review.....	104
Ten-year perspective	23	Medical Examiner activity.....	105
Manner of death: Accident.....	32	Organization of the King County Medical Examiner's Office in 2013	112
Manner of death: Homicide	41	Glossary of terms	113
Manner of death: Natural.....	48		
Manner of death: Suicide	58		



FOREWORD

The King County Medical Examiner's Office serves the community by investigating sudden, unexpected, violent, suspicious, and unnatural deaths. Medical Examiner staff recognize the tragedy surrounding an untimely death and perform investigations, in part, to assist the grieving family. A complete investigation provides for the quick settling of estates and insurance claims, as well as for implementing civil and criminal actions. Questions that seem irrelevant in the initial hours after death can become significant in the following months. The surviving family, friends, and general public can have the assurance that the Medical Examiner conducted a comprehensive investigation.

When a death occurs on the job or is work-related, the King County Medical Examiner's Office immediately forwards the results of its investigation to the Washington State Department of Labor and Industries so that the family can gain the full benefit of the findings. Private insurance companies also routinely use the findings to settle claims. Whenever a consumer product is implicated in a death, the King County Medical Examiner's Office notifies the Consumer Product Safety Commission to ensure that the product is studied and the necessary steps are taken to protect the public.

The public health role of the Medical Examiner is to isolate and identify the causes of sudden,

unexpected death that might affect more than one person. When an infectious agent or toxin is implicated in a death, the Medical Examiner's Office notifies the family and contacts of the deceased so they may receive any needed medical treatment. Trends in injury and violence are monitored. In this era of concern about emergency response and bioterrorism, the Medical Examiner provides an important level of preparedness and surveillance.

Civil or criminal judicial proceedings frequently require the medical investigation of violent death. Thus, the King County Medical Examiner's Office conducts a prompt medical investigation to provide the criminal justice system with medical information and evidence required for adjudication. Although criminal death investigations constitute a small portion of deaths investigated by the Medical Examiner, these deaths are studied in great detail because of the issues and legal consequences involved. The King County Medical Examiner's Office provides the criminal justice system the best support that medical science can provide.

In summary, the King County Medical Examiner's Office provides expert medical evaluation and extensive services related to the investigation of deaths that are of concern to the health, safety, and welfare of the community.



EXECUTIVE SUMMARY

The Medical Examiner's Office 2013 Annual Report reflects the activities pertaining to investigation of jurisdictional deaths in King County. The mission of the King County Medical Examiner's Office (KCMEO) is to investigate sudden, unexpected and unnatural deaths in King County with the highest level of professionalism, compassion and efficiency, and to provide a resource for improving the health and safety of the community.

This annual report presents detailed analyses of the different manners of deaths, as well as trends in homicides, traffic fatalities, and drug overdose deaths. While the report tends to depict the more violent types of deaths, it is worth noting that nearly 44% of Medical Examiner cases were classified as natural deaths.

In addition, data provided within this report helps shape Public Health policies designed to save lives by reducing preventable deaths. This report also documents the Medical Examiner's role in support of life saving organ and tissue donations, see page 103 for further details.

A few selected findings are highlighted below:

- In 2013, there were an estimated 13,639 deaths in King County. Of those deaths, 11,459 (84%) were reported to the Medical Examiner's Office. Deaths occurring in a hospital setting from a known natural disease process are not required to be reported to the Medical Examiner's Office. The Medical Examiner's Office assumed jurisdiction over 2,205 deaths; the number of applicable cases used in this report is 2,119 deaths after non-human remains and contract anthropology cases for other jurisdictions are removed. The King County Medical Examiner's Office assumes jurisdiction if a death falls under the Revised Code of Washington that defines the Medical Examiner's charge.
- The Medical Examiner's Office performed autopsies in 63% of those jurisdictional deaths (1,326/2,119). In 2013, those jurisdictional deaths included: 74 homicides, 266 suicides, 123 traffic deaths, 668 accidental deaths, 922 natural deaths and 66 deaths due to undetermined causes.
- Of the 22 natural deaths of children and youth investigated by the Medical Examiner, 59% (13/22) were of infants less than one year of age. Of those 13 infants who died of natural causes, 7 were due to Sudden Infant Death Syndrome (SIDS). In addition, 10 infant deaths were classified as "Sudden Unexplained Infant Death" (SUID), manner undetermined, due to the inability to exclude external factors that might have contributed to the death.
- Several factors appear repeatedly in unnatural deaths. Of all traffic fatalities in which tests were performed, 21% tested positive for the presence of alcohol in the blood. Firearms were the most frequent instrument of death in homicides (59%) and suicides (38%).

- Males comprised 82% (61/74) and women 18% (13/74) of the homicide victims in 2013. The majority of victims, 62% (46/74), were between the age 20 and 49. The number of homicide victims 19 years old and under remained the same when compared to the previous year. In 2013 they accounted for 15% (11/74) of the homicide victims, compared to 2012 when this younger age group represented 16% (11/69) of all homicide victims. 88% (65/74) of the victims were tested for the presence of alcohol. Of those tested 32% (21/65) showed alcohol present at the time of death.
- In 2013, there were 44 firearm homicide victims, 16% (7/44) were 19 years old and younger – a percentage increase from 2012 when 15% (7/47) of firearm homicide victims were 19 years old and younger. In 2013, there was a disproportionate number of firearm homicide victims that were African American (48%, 21/44) when compared to the percentage of African Americans in King County's population (6.3%) (see discussions on pages 8 and 44.) Of the 21 African American firearm homicide victims, 52% (11/21) were males 29 years old and younger. In comparison, 39% (17/44) of all the homicide firearm victims were White. Of the 17 White firearm homicide victims, 59% (10/17) were males 29 years old and younger.
- For King County in 2013, drugs and poisons caused 329 deaths, approximately 16% of all deaths investigated (329/2,119). The total number of drug-caused deaths increased compared to 2012 when there were 298 drug deaths. In 2013, deaths due to drugs and poisons comprised 33% (329/1000) of all suicidal, accidental and undetermined deaths combined.
- In 2013 the King County Medical Examiner's Office maintained accreditation by the National Association of Medical Examiners. This is the national professional organization of physician medical examiners, medicolegal death investigators and death investigation system administrators who perform the official duties of the medicolegal investigation of deaths in the United States.

Description and purpose

In 1969, the King County Home Rule Charter abolished the King County Office of the Coroner, which was replaced with the King County Medical Examiner's Office. The Medical Examiner's Office is a part of the Prevention Division of Public Health – Seattle & King County. The King County Medical Examiner's Office is funded by King County and operates under the direction of the King County Executive.

The Chief Medical Examiner, Dr. Richard Harruff, is a physician trained and certified in forensic pathology - the branch of medicine devoted to the scientific investigation of sudden, unexpected, violent, suspicious, or unnatural deaths. There are four sections under the Chief Medical Examiner's direction: Forensic Pathology, Scene Investigation, Autopsy Support and Administrative Support. The duties of these four sections include the performance of autopsies, certification of death, field investigation of scene and circumstances of death, identification of the deceased, notification of next-of-kin, and control and disposition of the deceased's personal property.

Deaths that come under the jurisdiction of the Medical Examiner are defined by state statute (RCW 68.50) and include, but are not limited to, the following circumstances:

1. *Persons who die suddenly when in apparent good health and without medical attendance within 36 hours preceding death.* This category is reserved for the following situations: (1) Sudden death of an individual with no known natural cause for the death. (2) Death during an acute or unexplained rapidly fatal illness, for which a reasonable natural cause has not been established. (3) Death of an individual who was not under the care of a physician. (4) Death of a person in a nursing home or care facility where medical treatment is not provided by a licensed physician.
2. *Circumstances which indicate death was caused in part or entirely by unnatural or unlawful means.* This category includes but is not limited to: (1) Drowning, suffocation, smothering, burns, electrocution, lightning, radiation, chemical or thermal injury, starvation, environmental exposure, or neglect. (2) Unexpected death during, associated with, or as a result of diagnostic or therapeutic procedures. (3) All deaths in the operating room whether due to surgical or anesthetic procedures. (4) Narcotics or other drugs including alcohol or toxic agents, or toxic exposure. (5) Death of the mother caused by known or suspected abortion. (6) Death from apparent natural causes during the course of a criminal act, e.g., a victim collapses during a robbery. (7) Death that occurs within one year following an accident, even if the accident is not thought to have contributed to the cause of death. (8) Death following all injury-producing accidents, if recovery was considered incomplete or if the accident is thought to have contributed to the cause of death (regardless of the interval between the accident and death).
3. *Suspicious circumstances.* This category includes, but is not limited to, deaths under the following circumstances: (1) Deaths resulting from apparent homicide or suicide. (2) Hanging, gunshot wounds, stab wounds, cuts, strangulation, etc. (3) Alleged rape, carnal knowledge, or sodomy. (4) Death during the course of, or precipitated by, a criminal act. (5) Death that occurs while in a jail or prison, or while in custody of law enforcement or other non-medical public institutions.

4. *Unknown or obscure causes.* This category includes: (1) Bodies that are found dead. (2) Death during or following an unexplained coma.
5. *Deaths caused by any violence whatsoever, when the injury was the primary cause or a contributory factor in the death.* This category includes, but is not limited to: (1) Injury of any type, including falls. (2) Any death due to or contributed to by any type of physical trauma.
6. *Contagious disease.* This category includes only those deaths wherein the diagnosis is undetermined and the suspected cause of death is a contagious disease which may be a public health hazard.
7. *Unclaimed bodies.* *This category is limited to deaths where no next of kin or other legally responsible representatives can be identified for disposition of the body.*
8. *Premature and stillborn infants.* *This category includes only those stillborn or premature infants whose birth was precipitated by maternal injury or drug use, criminal or medical negligence, or abortion under unlawful circumstances.*

Mission Statement

The mission of the King County Medical Examiner's Office (KCMEO) is to investigate sudden, unexpected and unnatural deaths in King County with the highest level of professionalism, compassion and efficiency and to provide a resource for improving the health and safety of the community consistent with the general mission of Public Health.

To achieve this mission, the KCMEO will:

- Coordinate investigative efforts with law enforcement, hospitals, and other agencies in a professional and courteous manner.
- Treat decedents and their effects with dignity and respect, and without discrimination.
- Conduct investigations and autopsies professionally, scientifically, and conscientiously; and complete reports expeditiously with regard for the concerns of family members, criminal justice, and public health and safety.
- Provide compassion, courtesy, and honest information to family members and, with sensitivity for cultural differences, make appropriate efforts in assisting with their grief, medical and legal questions, disposition of decedents and effects, and other settlements.
- Collect, compile, and disseminate information regarding deaths in a manner consistent with the laws of Washington state and consistent with the mission of Public Health.
- Provide medical and scientific testimony in court and in deposition as well as medicolegal consultation for prosecuting attorneys, defense attorneys, and attorneys representing surviving family members.
- Promote and advance, through education and research, the sciences and practices of death investigation, pathology, and anthropology within KCMEO and in collaboration with educational institutions.
- Promote and maintain an emotionally and physically healthy and safe working environment for KCMEO employees, following Public Health policies for standards of conduct, management, and support for employee diversity, training, and development.
- Expand communication throughout Public Health and the community at large regarding the roles, responsibilities, and objectives of KCMEO.

Explanation of data

The Medical Examiner serves the geographic area that includes all 2,130 square miles of King County, bounded by Pierce County to the south, Snohomish County to the north, Kittitas and Chelan Counties to the east, and Puget Sound to the west. In 2010, the King County population was estimated to be 1,942,600.¹ Included within King County are 39 cities and towns including Seattle, the state's largest city. Mercer Island, Vashon Island, two major airports and several colleges and universities are all in the geographic area served by the Medical Examiner's Office. In King County there are more than 20 hospitals and one major trauma center which serves the entire Pacific Northwest region.

The KCMEO assumes jurisdiction of deaths occurring in King County that include both King County residents and nonresidents. King County residents who die in other counties do not fall under KCMEO jurisdiction. For data on deaths of King County residents, along with other health indicators, please see Public Health—Seattle & King County Community Health Indicators online at: www.kingcounty.gov/healthservices/health/data/chi.

This report summarizes demographics from individual cases in which the Medical Examiner assumed jurisdiction and presents them in aggregate form. Table 1-8 (Nearest Incorporated City to the Fatal Incident) on pages 21 and 22 represents the location of the incident to the nearest city, not the residential address of the individual. Each manner (category) of death is subdivided into the various sub-groupings (methods) appropriate to that manner, which together form a more detailed description of the cause and manner of death.

The variables displayed in the tables such as race, gender, age, etc., have been selected as those most likely to assist and interest individuals using this data in assembling a profile of statistics on deaths examined by the Medical Examiner's Office for 2013. The Washington State Office of Financial Management estimates the racial distribution of King County to be 74.7% White, 6.3% African American, 3.6% Two or More Races, 14.4% Asian/Pacific Islander (including Hawaiian and other Pacific Islanders), and 1% American Indian/Alaska Native.² Information on Hispanic ethnicity of the decedent is not available for every case, and will not be presented in this report.

Medical Examiner figures cannot be directly compared to the racial distribution of King County residents. This is because as mentioned above and emphasized in Table 1-9 on page 22, in 13% of the Medical Examiner cases the incident leading to death occurred outside of King County and the decedent likely was not a resident of King County. However, as a rough estimate, the only manner of death that varies from the racial distribution of the county by a large percentage is Homicide (see discussion on page 44).

Age groups displayed in the tables are divided into youth and adult. The youth groups are infants (newborn to 11 months), toddlers (1-5 years), grade school (6-12 years), junior high (13-15 years), and high school (16-19 years). Adult age groups are in corresponding decades with the last being 90 years of age or older.

¹State of Washington, Office of Financial Management, 2012 estimate.

² State of Washington, Office of Financial Management, 2010 estimate. (latest figures available)

Blood alcohol (ethanol) data included here represent the blood level at the time of death. Alcohol is metabolized at a rate of 0.015 to 0.018 grams percent per hour. Thus, if there is a significant survival interval, the blood alcohol at the time of death will be lower than at the time of incident. Consequently, blood alcohol tests are not performed in cases where death occurs more than 24 hours after the fatal injury. For these reasons, an unknown number of cases not tested or showing no blood alcohol may actually have had a measurable alcohol concentration at the time of incident.

Three sections are included that review specific issues: deaths due to drugs, deaths due to firearms, and deaths among children and youth. The firearm data pertain to the victim because data relating to the shooter are not included in the Medical Examiner's investigation. For deaths among children and youth, the analysis focuses on violent, non-natural causes of death.

Data on natural deaths is included. However, these deaths due to natural causes are not representative of all natural deaths in King County. Natural deaths that the Medical Examiner investigates are those that occur suddenly and unexpectedly with no physician in attendance, or under suspicious circumstances. Such natural deaths comprised 43% (922/2,119) of all deaths that the Medical Examiner's Office investigated in 2013.

The "undetermined" category includes deaths in which the manner could not be clearly determined. In some cases, serious doubt existed as to whether the injury occurred with intent or as a result of an accident. In others, lack of witnesses or prolonged time between death and discovery precluded the accurate determination of the circumstances surrounding death. Moreover, it may be difficult to assess street drug or medication overdose deaths as showing enough features to reasonably determine the manner of death. Also included in the undetermined category are fetal deaths, which, according to the State of Washington death certification guidelines, are not assigned a manner of death.

Medical Examiner cases in 2013

The following provides a summary of the raw data from the Medical Examiner's cases for the year 2013. Ten-year trends are shown beginning on page 23.

In 2013, there were an estimated 13,639 deaths that occurred in King County (0.70% of a 2010 population estimate of 1,942,600).³ A total of 11,459 (84%, 11,459/13,639) were reported to the Medical Examiner's Office by medical and law enforcement personnel. Based on analysis of the scene and circumstances of death and the decedent's medical history gathered by the forensic medicolegal death investigators, the Medical Examiner's Office assumed jurisdiction in 2,205 of these reported deaths, of which 86 were either ultimately found to be non-human remains or contract cases (i.e., cases in which autopsy and/or anthropology cases are examined for other counties or agencies). Throughout the report, except where stated, the non-human, anthropology, and contract cases are excluded. Thus, the Medical Examiner assumed jurisdiction in 16% (2,119/13,639) of deaths that occurred in King County in 2013.⁴

In approximately 82% (9,340/11,459) of the reported deaths, the Medical Examiner did not assume jurisdiction and perform an investigation; instead a "No Jurisdiction Assumed" (NJA) number was assigned. In such instances a physician with knowledge and awareness of the decedent's state of health certified the death. These are primarily natural deaths, with a predominance of individuals in nursing homes with a known fatal disease process. Of note is the fact that the Medical Examiner declined jurisdiction in 9,340 of the deaths that were reported. The Medical Examiner's Office applies a strict interpretation of its governing legislative language "persons who die suddenly when in apparent good health and without medical attendance within thirty-six hours preceding death" (RCW 68.50). The Medical Examiner assumes jurisdiction only if both conditions (lack of medical care and apparent good health) apply, and there is no attending outside physician with sufficient knowledge of the individual's natural disease condition to certify the death.

The Medical Examiner's Office performed autopsies in 63% (1,326/2,119) of the cases in which jurisdiction was assumed. Autopsies by a Medical Examiner pathologist were not performed in deaths where scene, circumstances, medical history, and external examination of the body provided sufficient information for death certification. In 2013, there were 335 such deaths, accounting for 16% (335/2,119) of the total deaths. In addition, there were 444 deaths, (21% 444/2,119) certified by attending private physicians after review by and consultation with the Medical Examiner.

Of all the traffic fatalities in which tests were performed, 29% (26/91) tested positive for presence of alcohol (ethanol) in the blood. In recognition of the importance of safety devices in traffic accidents, Medical Examiner data indicate that of the 68 vehicle occupants who died, 51% (35/68) were wearing seatbelt restraints.

In the 22 deaths involving motorcyclists, 16 (73% 16/22) were wearing helmets.

³Death certificates filed in King County, Vital Statistics, Public Health - Seattle & King County, March, 2012.

⁴Does not include non-human remains or anthropology/contract cases.

Firearms were the most frequent instrument of death in homicides and suicides, accounting for 59% (44/74) of the homicides and 38% (100/266) of the suicides.

While the discussion here tends to depict the more violent types of death, the reader should be reminded that 43% (922/2,119) of Medical Examiner cases involve natural deaths. Specific discussion and presentation of relevant tables regarding 2013 cases follow this brief summary.

Table 1-1 Deaths Occurring in King County / Medical Examiner Cases / 2013

CASES BY MANNER OF DEATH ⁵	NUMBER OF KCME DEATHS	PERCENT OF KCME DEATHS
Accident Other (A)	668	32%
Accident Traffic (T)	123	6%
Homicide (H)	74	3%
Natural (N)	922	43%
Suicide (S)	266	13%
Undetermined ⁶ (U)	66	3%
Total KCME general cases	2,119	
Non-applicable cases where jurisdiction was assumed	86	
Total KCME jurisdiction cases	2,205	
Total KCME general cases ⁷	2,119	
Deaths reported to KCME but no jurisdiction was assumed (NJA)	9,340	
All other deaths in King County not reported to KCME	2,180	
ALL KING COUNTY DEATHS ⁸	13,639	

⁵The letters following each manner of death will be used in most tables throughout this report.


⁶Includes four fetal deaths, which according to Washington State death certification procedures, are not assigned a manner of death.

⁷This is the total number of cases that will be referred to throughout this report unless otherwise noted.


⁸Death certificates filed in King County, Vital Statistics, Public Health - Seattle & King County, May 2013

Graph 1-1 All King County Deaths with Medical Examiner Jurisdiction Shown / 2013

Total Deaths in King County, 2013: 13,639



Graph 1-2 Manner of Death for All Medical Examiner Jurisdiction Cases / 2013
 Jurisdiction assumed in 2,119 cases.⁹



⁹This number does not include 86 non-applicable cases (non-human tissue/bones and anthropology/contract cases).

Graph 1-3 Method of Certification for all King County Medical Examiner Jurisdiction Cases / 2013




Table 1-2 Method of Certification / Manner of Death / KCME / 2013

CERTIFICATION	MANNER OF DEATH						TOTAL	%
	A	T	H	N	S	U		
KCME Autopsies	388	85	70	477	243	63	1,326	62.5%
KCME External Exams	98	28	0	184	23	2	335	15.8%
KCME Other	7	3	4	0	0	0	14	0.7%
Attending Physician	175	7	0	261	0	1	444	21%
Totals	668	123	74	922	266	66	2,119	100%

Manner of Death in 2013

King County Medical Examiner's Office General Cases

Graph 1-4 Gender / Manner of Death / KCME / 2013




Table 1-3 Gender / Manner of Death / KCME / 2013

GENDER	MANNER OF DEATH						TOTAL	%
	A	T	H	N	S	U		
Male	409	89	61	631	197	46	1,433	68%
Female	259	34	13	291	69	20	686	32%
Totals	668	123	74	922	266	66	2,119	100%

Table 1-4 Age / Gender / Manner of Death / KCME / 2013

AGE / GENDER	MANNER OF DEATH						Sub-Total	TOTAL	%
	A	T	H	N	S	U			
Under 1 year	1	0	2	13	0	17		33	1.6%
<i>Male</i>	1	0	1	9	0	10	21		
<i>Female</i>	0	0	1	4	0	7	12		
1-5 years	3	1	1	7	0	1		13	0.6%
<i>Male</i>	1	1	0	6	0	1	9		
<i>Female</i>	2	0	1	1	0	0	4		
6-12 years	1	0	0	1	2	0		4	0.2%
<i>Male</i>	1	0	0	0	2	0	3		
<i>Female</i>	0	0	0	1	0	0	1		
13-15 years	1	3	0	0	3	0		7	0.3%
<i>Male</i>	1	1	0	0	2	0	4		
<i>Female</i>	0	2	0	0	1	0	3		
16-19 years	5	9	8	3	10	1		36	1.7%
<i>Male</i>	5	7	5	3	9	1	30		
<i>Female</i>	0	2	3	0	1	0	6		
20-29 years	52	25	26	19	39	5		166	7.8%
<i>Male</i>	39	17	24	12	28	5	125		
<i>Female</i>	13	8	2	7	11	0	41		
30-39 years	68	6	14	41	45	8		182	8.6%
<i>Male</i>	51	6	12	24	37	5	135		
<i>Female</i>	17	0	2	17	8	3	47		
40-49 years	90	15	6	104	36	14		265	12.5%
<i>Male</i>	64	14	5	78	22	11	194		
<i>Female</i>	26	1	1	26	14	3	71		
50-59 years	112	20	9	210	47	10		408	19.3%
<i>Male</i>	71	15	8	165	34	9	302		
<i>Female</i>	41	5	1	45	13	1	106		
60-69 years	66	15	5	253	47	2		388	18.3%
<i>Male</i>	44	12	4	181	34	0	275		
<i>Female</i>	22	3	1	72	13	2	113		
70-79 years	62	13	1	150	17	0		243	11.5%
<i>Male</i>	39	8	1	97	12	0	157		
<i>Female</i>	23	5	0	53	5	0	86		
80-89 years	112	12	2	83	14	5		228	10.7%
<i>Male</i>	53	6	1	40	11	2	113		
<i>Female</i>	59	6	1	43	3	3	115		
90+years	95	4	0	38	6	3		146	6.9%
<i>Male</i>	40	2	0	11	6	2	61		
<i>Female</i>	55	2	0	27	0	1	85		
Totals	668	123	74	922	266	66		2,119	100%

Table 1-5 Race / Gender / Manner of Death / KCME / 2013¹⁰

RACE / GENDER	MANNER OF DEATH						Sub-Total	TOTAL	%
	A	T	H	N	S	U			
White	567	101	35	770	225	42		1,740	82%
<i>Male</i>	337	71	29	524	166	28	1,115		
<i>Female</i>	230	30	6	246	59	14	585		
African American	46	10	28	83	11	9		187	9%
<i>Male</i>	32	8	25	60	8	7	140		
<i>Female</i>	14	2	3	23	3	2	97		
Asian/Pacific Is.	40	7	5	56	26	6		140	7%
<i>Male</i>	30	5	4	37	19	5	100		
<i>Female</i>	10	2	1	19	7	1	40		
American Indian / Alaska Native	15	5	5	13	3	9		50	2%
<i>Male</i>	10	5	3	10	3	6	37		
<i>Female</i>	5	0	2	3	0	3	13		
Other	0	0	1	0	1	0		2	<0.1%
<i>Male</i>	0	0	0	0	1	0	1		
<i>Female</i>	0	0	1	0	0	0	1		
Totals	668	123	74	922	266	66		2,119	100%

¹⁰ A = Accident (Non-Traffic), T = Traffic, H = Homicide, N = Natural, S = Suicide, U = Undetermined.




Table 1-6 Marital Status / Gender / Manner of Death / KCME / 2013¹¹

MARITAL STATUS / GENDER	MANNER OF DEATH						Sub-Total	TOTAL	%
	A	T	H	N	S	U			
Never Married	208	58	48	324	117	44		799	37.7%
<i>Male</i>	160	46	40	254	89	32	621		
<i>Female</i>	48	12	8	70	28	12	178		
Married	174	39	11	177	78	9		488	23%
<i>Male</i>	115	29	10	117	59	7	337		
<i>Female</i>	59	10	1	60	19	2	151		
Divorced	133	14	10	251	60	8		476	22.5%
<i>Male</i>	76	10	8	170	41	6	311		
<i>Female</i>	57	4	2	81	19	2	165		
Widowed	137	10	4	110	8	4		273	12.9%
<i>Male</i>	46	3	2	40	5	0	96		
<i>Female</i>	91	7	2	70	3	4	177		
Unknown	15	1	1	59	3	1		80	3.8%
<i>Male</i>	11	1	1	49	3	1	66		
<i>Female</i>	4	0	0	10	0	0	14		
Domestic Partner	1	1	0	1	0	0		3	0.1%
<i>Male</i>	1	0	0	1	0	0	2		
<i>Female</i>	0	1	0	0	0	0	1		
Totals	668	123	74	922	266	66		2119	100%

¹¹A = Accident (Non-Traffic), T = Traffic, H = Homicide, N = Natural, S = Suicide, U = Undetermined.

Graph 1-6 Month / Manner of Death / KCME / 2013




Table 1-7 Month / Manner of Death / KCME / 2013¹²

MONTH	MANNER OF DEATH						Total	%
	A	T	H	N	S	U		
Prior to 2012	0	0	0	0	2	0	2	0.1%
2012	4	0	0	8	1	2	15	0.7%
January	70	14	3	79	17	7	190	9%
February	50	6	2	78	24	4	164	7.7%
March	56	7	12	94	19	6	194	9.2%
April	54	5	9	72	20	5	165	7.8%
May	55	15	11	72	25	3	181	8.5%
June	46	4	7	82	17	5	161	7.6%
July	46	13	6	73	26	4	168	7.9%
August	65	10	4	62	18	7	166	7.8%
September	59	15	6	79	29	3	191	9%
October	56	13	6	81	17	7	180	8.5%
November	55	11	3	75	30	8	182	8.6%
December	52	10	5	67	21	5	160	7.6%
Totals	668	123	74	922	266	66	2,119	100%

¹²A = Accident (Non-Traffic), T = Traffic, H = Homicide, N = Natural, S = Suicide, U = Undetermined.

Table 1-8 Nearest Incorporated City to the Fatal Incident / KCME / 2013¹³

CITY	MANNER OF DEATH					TOTAL	%
	A	T	H	S	U		
Algona	1	0	0	0	0	1	0.1%
Auburn	31	5	4	11	4	55	4.6%
Beaux Arts	0	0	0	0	0	0	0%
Bellevue	41	3	1	10	0	55	4.6%
Black Diamond	0	0	0	0	0	0	0%
Bothell	10	0	0	3	1	14	1.2%
Burien	10	1	0	8	1	20	1.7%
Carnation	1	0	0	1	0	2	0.2%
Clyde Hill	0	0	0	0	0	0	0%
Covington	1	0	1	2	0	4	0.3%
Des Moines	11	0	0	5	1	17	1.4%
Duvall	1	0	0	3	0	4	0.3%
Enumclaw	6	2	1	3	0	12	1%
Federal Way	25	4	6	15	2	52	4.3%
Hunts Point	0	0	0	0	0	0	0%
Issaquah	7	3	0	4	0	14	1.2%
Kenmore	3	2	0	6	0	11	0.9%
Kent	26	8	6	15	3	58	4.9%
Kirkland	20	2	0	6	3	31	2.6%
Lake Forest Park	2	0	0	3	0	5	0.4%
Maple Valley	4	1	2	1	1	9	0.8%
Medina	0	0	0	0	1	0	0.1%
Mercer Island	4	0	0	0	0	4	0.3%
Milton	0	0	0	0	0	0	0%
Newcastle	1	0	0	0	0	1	0.1%
Normandy Park	1	0	0	2	0	3	0.3%
North Bend	8	1	1	6	1	17	1.4%
Pacific	2	0	0	2	0	4	0.3%

¹³ Table does not include cases where manner of death is classified "Natural". A = Accident (Non-Traffic), T = Traffic, H = Homicide, S = Suicide, U = Undetermined.

Table 1-8 Nearest Incorporated City to the Fatal Incident / KCME / 2013¹⁴ (continued)

CITY	MANNER OF DEATH					Total	%
	A	T	H	S	U		
Redmond	16	3	0	8	1	28	2.3%
Renton	27	4	6	10	1	48	4%
Sammamish	2	0	0	5	0	7	0.6%
SeaTac	7	1	2	3	0	13	1.1%
Seattle	278	34	32	97	33	475	39.7%
Shoreline	12	3	0	8	1	24	2%
Skykomish	2	0	0	1	0	3	0.2%
Snoqualmie	1	1	0	1	1	4	0.3%
Tukwila	6	0	1	2	0	9	0.8%
Woodinville	7	0	1	6	0	14	1.2%
Yarrow Point	0	0	0	0	0	0	0%
Unincorporated King County							
Hobart	1	0	0	0	0	1	0.1%
Fall City	0	0	0	1	0	1	0.1%
Preston	0	1	0	0	0	1	0.1%
Ravensdale	1	0	0	2	0	3	0.2%
Vashon Island	0	1	0	6	0	7	0.6%
Outside of King County	88	41	10	8	8	157	13%
Unknown Location	4	2	0	0	2	8	0.7%
Totals	668	123	74	266	66	1197	100%

¹⁴A = Accident (Non-Traffic), T = Traffic, H = Homicide, S = Suicide, U = Undetermined.

Out of County Cases 2013

King County is home to many hospitals and a major trauma center that serves the entire Pacific Northwest and the western United States. Consequently, there are numerous deaths each year where the incident leading to death occurred outside of King County. However, because the death occurred within King County, it comes under the jurisdiction of the King County Medical Examiner's Office. In 2013, there were 163 deaths (14%, 163/1,197) where the incident (excluding deaths classified as "Natural") occurred out of county or where the incident location was unknown. Table 1-9 displays these deaths by incident location and manner.

Table 1-9 Fatal Incident Occurred Outside of King County / KCME / 2013¹⁵

INCIDENT LOCATION	MANNER OF DEATH					TOTAL
	A	T	H	S	U	
Alaska	1	0	0	1	1	3
Montana	4	0	0	0	0	4
Idaho	1	3	1	0	0	5
Oregon	1	2	1	0	0	4
Other States	0	2	0	0	0	2
Washington						
<i>Island County</i>	3	3	0	0	2	8
<i>Kitsap County</i>	10	0	0	0	0	10
<i>Pierce County</i>	5	1	0	1	1	8
<i>Skagit County</i>	8	2	0	0	0	10
<i>Snohomish County</i>	22	12	4	3	2	43
<i>Thurston County</i>	0	2	0	3	0	5
<i>Other WA Counties</i>	33	14	4	2	2	55
Washington Sub-Total	81	34	8	9	7	139
Out of Country	0	0	0	0	0	0
Unknown	4	0	0	0	2	6
Totals	92	41	10	10	10	163

¹⁵Table does not include cases where manner of death is classified as "Natural." A = Accident (Non-Traffic), T = Traffic, H = Homicide, S = Suicide, U = Undetermined.

Ten-year perspective

This section provides a ten-year perspective on deaths that the Medical Examiner investigated and shows the variations in data from year to year.

The tables on the following pages attempt to give a perspective on the types of deaths that the Medical Examiner investigates. The tables display data by category and year and provide trends over time. More detailed analysis of 2013 data is provided in separate sections for each manner of death (Accident, Homicide, Natural, Suicide, Traffic, and Undetermined).


Table 2-1 Comparison of Manners of Death / KCME / 2004 - 2013

MANNER OF DEATH	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Accident (Other)	542	602	721	687	739	632	593	594	670	668
Accident (Traffic)	192	226	211	170	163	141	150	135	131	123
Homicide	76	80	91	76	85	63	59	54	69	74
Natural	765	763	752	863	871	989	953	926	873	922
Suicide	229	233	227	223	210	253	232	265	281	266
Undetermined	62	41	53	53	53	59	73	62	80	66
Totals	1,866	1,945	2,055	2,072	2,121	2,137	2,060	2,036	2,104	2,119

Table 2-2 Comparison of Manners of Death as Percentage of Total Annual Medical Examiner Cases / KCME / 2004 – 2013

MANNER OF DEATH	2004	2005	2006	2007	2008	2009	2010	2010	2012	2013
	%	%	%	%	%	%	%	%	%	%
Accident (Other)	29.0	31.0	35.1	33.1	34.8	29.6	28.8	29.2	31.8	31.5
Accident (Traffic)	10.3	11.6	10.3	8.2	7.7	6.6	7.3	6.6	6.2	5.8
Homicide	4.1	4.1	4.4	3.7	4.0	2.9	2.9	2.7	3.3	3.5
Natural	41.0	39.2	36.6	41.7	41.1	46.3	46.3	45.5	41.5	43.5
Suicide	12.3	12.0	11.0	10.8	9.9	11.8	11.2	13	13.4	12.6
Undetermined	3.3	2.1	2.6	2.5	2.5	2.8	3.5	3	3.8	3.1
Totals	100%									

Graph 2-1 Comparison of Manners of Death / KCME / 2004- 2013



Graph 2-2 Homicide Deaths / KCME / 2004 - 2013




Table 2-3 Ten-Year Perspective of Homicidal Methods / KCME / 2004 – 2013

METHOD USED	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Blunt Force (#)	10	12	16	9	16	5	11	6	6	14
Blunt Force (%)	13%	15%	18%	12%	19%	8%	18%	11%	9%	19%
Firearms (#)	46	47	52	55	45	41	39	35	47	44
Firearms (%)	61%	59%	57%	72%	53%	65%	66%	65%	68%	59%
Hom. Violence (#)	3	2	0	0	0	0	1	1	3	0
Hom. Violence (%)	4%	3%	0%	0%	0%	0%	2%	2%	4%	0%
Stabbing (#)	10	14	14	12	12	11	2	9	13	11
Stabbing (%)	13%	17%	15%	16%	14%	17%	4%	16%	19%	15%
Strangulation (#)	1	4	1	0	4	3	1	2	0	3
Strangulation (%)	1%	5%	1%	0%	5%	5%	2%	4%	0%	4%
Other (#)	6	1	8	0	8	3	5	1	0	2
Other (%)	8%	1%	9%	0%	9%	5%	8%	2%	0%	3%
Totals	76	80	91	76	85	63	59	54	69	74

Graph 2-3 Suicide Deaths /KCME / 2004 – 2013




Table 2-4 Ten Year Perspective of Suicidal Injury Modes / KCME / 2004 - 2013

INJURY MODE	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Asphyxia / Plastic Bag	7	5	11	3	8	8	13	15	21	13
Burns / Fire	1	3	3	1	3	2	2	1	2	1
Carbon Monoxide	8	13	11	17	4	14	4	7	9	10
Drowning	5	0	1	3	3	7	3	5	7	2
Drugs / Poisons	41	39	36	36	29	29	43	41	42	41
Firearms	95	96	98	93	93	100	92	116	119	100
Hanging	44	42	31	43	48	60	44	48	48	71
Incised Wounds / Stabbing	8	9	5	4	5	8	7	12	8	9
Jumped	15	22	26	22	13	20	21	19	24	15
Other	5	4	5	1	4	5	3	1	1	4
Totals	229	233	227	223	210	253	232	265	281	266

Graph 2-4 Traffic Fatalities / KCME / 2004 – 2013





Table 2-5 Traffic Fatality Circumstances / KCME / 2004 - 2013

CIRCUMSTANCES	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Vehicle Driver	78	99	92	71	71	51	69	55	47	45
Vehicle Passenger	54	47	44	29	24	28	27	22	16	23
Vehicle Unknown Position	1	1	5	1	4	0	0	3	4	0
Bicyclist	5	6	8	7	4	12	3	8	5	7
Motorcycle Driver	23	33	27	26	28	18	24	26	24	22
Motorcycle Passenger	0	3	1	2	1	1	0	1	1	0
Pedestrian	30	36	33	31	26	29	27	17	33	25
Other	1	1	1	3	5	2	0	3	1	1
Totals	192	226	211	170	163	141	150	135	131	123


Table 2-6 Ten Year Perspective of Non-Traffic Accidental Death Circumstances / KCME / 2004 - 2013

CIRCUMSTANCES	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Aircraft	2	3	3	11	1	0	0	1	3	1
Asphyxia	2	9	12	11	14	10	5	6	7	8
Aspiration	8	10	9	5	10	7	6	7	15	13
Blunt Force / Crushing	8	10	4	10	10	6	5	11	20	3
Burns / Fire	24	26	23	23	13	15	29	18	26	19
Carbon Monoxide	3	4	8	3	4	4	2	3	0	0
Drowning	17	19	30	23	23	17	11	21	24	23
Drugs / Poisons	211	216	262	247	232	233	214	203	230	279
Electrocution	2	1	2	1	1	2	2	1	1	2
Explosion	4	1	1	2	0	0	3	0	0	0
Fall	213	230	308	292	323	309	291	291	314	291
Firearms	1	2	0	1	1	1	1	0	2	1
Hanging	2	2	0	0	1	1	1	2	4	1
Hypothermia	2	4	4	3	4	7	4	7	6	5
Struck by Object	7	1	8	5	2	4	4	3	2	1
Struck by Train	3	1	0	1	3	2	0	6	2	5
Vehicular Non-Traffic	10	8	9	7	10	5	2	4	4	7
Other	5	10	7	2	6	9	13	10	10	9
Totals	542	602	721	687	739	632	593	594	670	668


Graph 2-5 Accidental Deaths / KCME / 2004 – 2013



Graph 2-6 Natural Deaths / KCME / 2004 – 2013



Graph 2-7 Deaths of Undetermined Manner / KCME / 2004 – 2013



Manner of death: Accident

The Medical Examiner certified 668 deaths as non-traffic accidents for the calendar year 2013. The largest group of accidental deaths was those who died as a result of a fall, 44% (291/668). Of the 291 deaths attributed to injury sustained in falls, 83% (242/291) occurred in the age group 70 years and over. The largest percentage was ground-level falls in elderly individuals, which resulted in fractures leading to complications such as pneumonia.

The second largest group of non-traffic accidental deaths was individuals who died as a result of accidental overdoses of drugs and/or poisons, representing 42% (279/668). There was one accidental drug death of a child between the ages of 16-19 years, and there was one death of a child less than 15 years of age.

The 2013 accidental drug death percentage, 42% (279/668) is eight percent greater than the 34% (227/670) accidental drug deaths in 2012. A more detailed discussion of these deaths is presented in the section "Death Due to Drugs and Poisons" on pages 89 and 90.

In 2013, 19 deaths resulted from fire or thermal injury, a decrease from 2012 when there were 26. Of the 19 fire-related deaths, 68% (13/19) were the result of accidents that occurred outside of King County. The injured were transported to Harborview Medical Center's Burn Intensive Care Unit where they died.

Another category of accidental deaths worthy of comment is death resulting from drowning. There were 23 drowning deaths in 2013, as compared to 24 in 2012.

Aspiration is a type of death that results from a person choking on a foreign object, often a bolus of food while eating. In 2013, there were thirteen deaths due to aspiration of a foreign body, compared to fifteen in 2012. All of the aspiration deaths were in adults over the age of 20.

Of the 668 accidental deaths in 2013, 14% (92/668) were the result of incidents which occurred outside of King County, but the death took place within King County. These deaths were the result of the injured being transported from outside King County to medical facilities within King County where they died. Since these deaths occurred in King County, they fall under King County Medical Examiner's Office jurisdiction.

Sixty percent (402/668) of the victims were tested for the presence of alcohol. Of those tested, 29% (118/402) showed alcohol present at the time of death.

Graph 3-1 Circumstances of Accidental Death / KCME / 2013





Table 3-1 Circumstances of Accidental Death / Race / Gender / KCME / 2013

CIRCUMSTANCES / GENDER	RACE					SUB-TOTAL	TOTAL
	WHITE	AFRICAN AMER	ASIAN/ PAC IS	AM INDIAN / AK NATIVE	OTHER		
Aircraft	1	0	0	0	0		1
<i>Male</i>	1	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	
Asphyxia: compressional / positional / mechanical	5	1	2	0	0		8
<i>Male</i>	5	1	1	0	0	7	
<i>Female</i>	0	0	1	0	0	1	
Aspiration	11	0	1	1	0		13
<i>Male</i>	4	0	1	1	0	6	
<i>Female</i>	7	0	0	0	0	7	
Blunt Force / Crushing	3	0	0	0	0		3
<i>Male</i>	3	0	0	0	0	3	
<i>Female</i>	0	0	0	0	0	0	
Burns / Fire	19	0	0	0	0		19
<i>Male</i>	11	0	0	0	0	11	
<i>Female</i>	8	0	0	0	0	8	
Complication of Therapy	1	0	0	0	0		1
<i>Male</i>	0	0	0	0	0	0	
<i>Female</i>	1	0	0	0	0	1	
Drowning	17	5	0	1	0		23
<i>Male</i>	11	5	0	1	0	17	
<i>Female</i>	6	0	0	0	0	6	
Drugs / Poisons	221	36	14	8	0		279
<i>Male</i>	144	24	13	6	0	187	
<i>Female</i>	77	12	1	2	0	92	
Electrocution	2	0	0	0	0		2
<i>Male</i>	2	0	0	0	0	2	
<i>Female</i>	0	0	0	0	0	0	

Table 3-1 Circumstances of Accidental Death / Race / Gender / KCME / 2013 (continued)

CIRCUMSTANCES / GENDER	RACE					SUB- TOTAL	TOTAL
	WHITE	AFRICAN AMER	ASIAN/ PAC IS	AM INDIAN/ AK NATIVE	OTHER		
Fall	264	2	21	4	0		291
<i>Male</i>	136	0	13	1	0	150	
<i>Female</i>	128	2	8	3	0	141	
Gunshot wound(s)	1	0	0	0	0		1
<i>Male</i>	1	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	
Hanging	1	0	0	0	0		1
<i>Male</i>	1	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	
Hypothermia	5	0	0	0	0		5
<i>Male</i>	2	0	0	0	0	2	
<i>Female</i>	3	0	0	0	0	3	
Non-Traffic Vehicular	5	1	1	0	0		7
<i>Male</i>	5	1	1	0	0	7	
<i>Female</i>	0	0	0	0	0	0	
Struck by Object	1	0	0	0	0		1
<i>Male</i>	1	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	
Struck by Train	4	0	0	1	0		5
<i>Male</i>	4	0	0	1	0	5	
<i>Female</i>	0	0	0	0	0	0	
Other	6	1	1	0	0		8
<i>Male</i>	6	1	1	0	0	8	
<i>Female</i>	0	0	0	0	0	0	
Totals	567	46	40	15	0		668
Percent	85%	7%	6%	2%	0%		100%

Graph 3-2 Accidental Deaths / Race / KCME / 2013



Graph 3-3 Accidental Deaths / Age Group / KCME / 2013




Table 3-2 Circumstances of Accidental Death / Age / Gender / KCME / 2013

CIRCUMSTANCES / GENDER	AGE GROUP (YEARS)													SUB-TOTAL	TOTAL
	< 1	1 to 5	6 to 12	13 to 15	16 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 to 69	70 to 79	80 to 89	90 +		
Aircraft	0	0	0	0	0	0	0	1	0	0	0	0	0	1	
Male	0	0	0	0	0	0	0	1	0	0	0	0	0	1	
Female	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Asphyxia compressional / positional / mechanical	1	0	1	0	0	0	1	1	2	1	1	0	0	8	
Male	1	0	1	0	0	0	1	1	1	1	1	0	0	7	
Female	0	0	0	0	0	0	0	0	1	0	0	0	0	1	
Aspiration	0	0	0	0	0	1	0	0	2	2	4	1	3	13	
Male	0	0	0	0	0	1	0	0	2	0	2	0	1	6	
Female	0	0	0	0	0	0	0	0	0	2	2	1	2	7	
Blunt Force / Crushing	0	0	0	0	1	1	0	0	0	0	1	0	0	3	
Male	0	0	0	0	1	1	0	0	0	0	1	0	0	3	
Female	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Burns / Fire	0	1	0	0	0	1	1	4	1	3	3	3	2	19	
Male	0	0	0	0	0	1	1	2	1	3	1	1	1	11	
Female	0	1	0	0	0	0	0	2	0	0	2	2	1	8	
Complication of Therapy	0	0	0	0	0	0	0	0	0	1	0	0	0	1	
Male	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Female	0	0	0	0	0	0	0	0	0	1	0	0	0	1	
Drowning	0	1	0	0	3	3	2	4	5	2	1	2	0	23	
Male	0	0	0	0	3	3	1	3	3	2	1	1	0	17	
Female	0	1	0	0	0	0	1	1	2	0	0	1	0	6	
Drugs / Poisons	0	0	0	1	1	41	58	69	76	29	4	0	0	279	
Male	0	0	0	1	1	28	45	47	42	19	4	0	0	187	
Female	0	0	0	0	0	13	13	22	34	10	0	0	0	92	
Electrocution	0	0	0	0	0	1	1	0	0	0	0	0	0	2	
Male	0	0	0	0	0	1	1	0	0	0	0	0	0	2	
Female	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Fall	0	1	0	0	0	1	3	7	16	21	47	105	90	291	
Male	0	1	0	0	0	1	0	6	13	13	28	50	38	150	
Female	0	0	0	0	0	0	3	1	3	8	19	55	52	141	
Gunshot wound(s)	0	0	0	0	0	0	0	1	0	0	0	0	0	1	
Male	0	0	0	0	0	0	0	1	0	0	0	0	0	1	
Female	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Hanging	0	0	0	0	0	0	0	1	0	0	0	0	0	1	
Male	0	0	0	0	0	0	0	1	0	0	0	0	0	1	
Female	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Hypothermia	0	0	0	0	0	0	0	0	2	2	1	0	0	5	
Male	0	0	0	0	0	0	0	0	1	1	0	0	0	2	
Female	0	0	0	0	0	0	0	0	1	1	1	0	0	3	

King County Medical Examiner's Office - 2013 Annual Report

Non-traffic Vehicular	0	0	0	0	0	0	1	1	4	1	0	0	0	7
<i>Male</i>	0	0	0	0	0	0	1	1	4	1	0	0	0	7
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Struck by Object	0	0	0	0	0	0	0	0	0	0	0	1	0	1
<i>Male</i>	0	0	0	0	0	0	0	0	0	0	0	1	0	1
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Struck by Train	0	0	0	0	0	2	0	0	2	1	0	0	0	5
<i>Male</i>	0	0	0	0	0	2	0	0	2	1	0	0	0	5
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Other	0	0	0	0	0	1	1	1	2	3	0	0	0	8
<i>Male</i>	0	0	0	0	0	1	1	1	2	3	0	0	0	8
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Totals	1	3	1	1	5	52	68	90	112	66	62	112	95	668
Percent	.1%	.5%	.1%	.1%	.7%	7.8%	10.2%	13%	16.8	9.9%	9.3%	16.8%	14.2%	100%

Table 3-3 Circumstances of Accidental Death / Gender / KCME / 2013

CIRCUMSTANCES	GENDER		TOTAL
	MALE	FEMALE	
Aircraft	1	0	1
Asphyxia (compressional / positional / mechanical)	7	1	8
Aspiration	6	7	13
Blunt Force / Crushing	3	0	3
Burns / Fire	11	8	19
Complication of Therapy	0	1	1
Drowning	17	6	23
Drugs / Poisons	187	92	279
Electrocution	2	0	2
Fall	150	141	291
Gunshot wound(s)	1	0	1
Hanging	1	0	1
Hypothermia	2	3	5
Non-traffic Vehicular	7	0	7
Struck by Object	1	0	1
Struck by Train	5	0	5
Other	8	0	8
Totals	409	259	668
Percent	61.2%	38.8%	100%

Table 3-4 Circumstances of Accidental Death / Blood Alcohol Results / KCME / 2013

CIRCUMSTANCES	TESTED		NOT TESTED	TOTAL
	TESTED POSITIVE	TESTED NEGATIVE		
Aircraft	0	1	0	1
Asphyxia (compressional/ positional / mechanical)	2	5	1	8
Aspiration	2	4	7	13
Blunt Force / Crushing	0	1	2	3
Burns / Fire	3	9	7	19
Complication of Therapy	0	1	0	1
Drowning	8	15	0	23
Drugs / Poisons	87	181	11	279
Electrocution	1	1	0	2
Fall	11	49	231	291
Gunshot wound(s)	0	1	0	1
Hanging	0	1	0	1
Hypothermia	0	5	0	5
Non-traffic Vehicular	0	5	2	7
Struck by Object	0	0	1	1
Struck by Train	3	2	0	5
Other	1	3	4	8
Totals	118	284	266	668
Percent	18%	42%	40%	100%

Manner of death: Homicide

The Medical Examiner classifies a death as a homicide when the death results from injuries inflicted by another person. In this context, the word homicide does not necessarily imply the existence of criminal intent behind the action of the other person. This is reflected in the fact that the prosecuting attorney may either charge the person responsible for the injuries with murder or manslaughter, or decline to file charges. In 2013, the Medical Examiner classified 74 deaths as homicide. This number represents 3.3% (74/2205) of the Medical Examiner death investigations for the calendar year 2013. Of these 74 homicides, 64 (86%, 64/74) were the result of incidents that occurred within King County. For comparison, there were 69 homicides investigated in 2012, of which 60 (87%, 60/69) were incidents in King County.

The data reflect the weapons or mechanisms responsible for the homicidal deaths in 2013. Firearms were responsible for 59% (44/74), compared to 2012, when 68% (47/69) were due to firearms. Stabbing by a knife or other sharp-edged instrument caused 15% (11/74) of deaths of homicide victims. Blunt force injuries were responsible for 19% (14/74) of the 2013 homicide deaths. There were three deaths due to strangulation/asphyxia, no deaths due to homicidal violence and two deaths due to other means. The term "homicidal violence" is used when circumstances indicate that death was due to homicide, but the exact cause of death is not determined, for example, in a decomposed body. There was one such death in 2012.

In 2013, there were three homicide victims under five years of age. There were no homicide victims between 6 - 15 years of age. Eight homicide victims were between the ages of 16 and 19 years.

Examining the racial distribution of victims of homicide, 38% (28/74) of the victims were African American, compared to 2012, when 33% (23/69) of the victims were African American. Whites, while representing 74.7% of the population, made up 47% (35/74) of the homicide victims. The remaining 15% of homicide victims (11/74) included Asian/Pacific Islanders (5/74) Native Americans/AK Natives (5/74) and other (1/74). As indicated on pages 9 and 23, in 14% of the Medical Examiner cases the incident leading to death occurred outside of King County and the decedent was likely not a resident of King County. Therefore, Medical Examiner figures cannot be directly compared to the racial distribution of King County residents (refer to Table 1-9 on page 23.)

Males comprised 82% (61/74) and women 18% (13/74) of the homicide victims in 2013. The majority of victims, 74% (55/74), were between the ages of 20 and 59 years. Young people, 19 years old and under, comprised 15% (11/74) of the homicide victims. For comparison, this younger age group represented 16% (11/69) in the year 2012. Eighty-eight percent (65/74) of the victims were tested for the presence of alcohol. Of those tested 32% (21/65) showed alcohol present at the time of death.

Of the 74 homicide deaths in 2013, 64(86% 64/74) of the fatal incidents occurred within King County, and of these deaths, 32 (50% 32/64) occurred within the city limits of Seattle. In 10 of the 74 homicidal deaths, the incident occurred outside of King County, but death occurred within King County.

The relationship of victim to assailant was not tabulated as part of this report. In order to investigate such associations, additional review of police records would be necessary.

Graph 4-1 Homicide Injury Methods / KCME / 2013




Table 4-1 Homicide Methods / Race / Gender / KCME / 2013


CIRCUMSTANCES / GENDER	RACE					SUB-TOTAL	TOTAL
	WHITE	AFRICAN AMER	ASIAN/ PAC IS	AM INDIAN/ AK NATIVE	OTHER		
Asphyxia / Strangulation	2	0	1	0	0		3
<i>Male</i>	1	0	0	0	0	1	
<i>Female</i>	1	0	1	0	0	2	
Blunt Force	8	3	0	3	0		14
<i>Male</i>	7	2	0	1	0	10	
<i>Female</i>	1	1	0	2	0	4	
Firearms	17	21	3	2	1		44
<i>Male</i>	16	20	3	2	0	41	
<i>Female</i>	1	1	0	0	1	3	
Homicidal Violence	0	0	0	0	0		0
<i>Male</i>	0	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	0	
Stabbing	6	4	1	0	0		11
<i>Male</i>	4	3	1	0	0	8	
<i>Female</i>	2	1	0	0	0	3	
Other / Unknown	2	0	0	0	0		2
<i>Male</i>	1	0	0	0	0	1	
<i>Female</i>	1	0	0	0	0	1	
Totals	35	28	5	5	1		74
Percent	47%	38%	7%	7%	1%		100%

Table 4-2 Homicide Methods / Age / Gender / KCME / 2013

AGE GROUP (YEARS)

METHOD / GENDER	< 1	1	6	13	16	20	30	40	50	60	70	80	90	SUB-TOTAL	TOTAL
		to 5	to 12	to 15	to 19	to 29	to 39	to 49	to 59	to 69	to 79	to 89	+		
Asphyxia / Strangulation	0	0	0	0	0	0	0	1	0	0	0	2	0		3
<i>Male</i>	0	0	0	0	0	0	0	0	0	0	0	1	0	1	
<i>Female</i>	0	0	0	0	0	0	0	1	0	0	0	1	0	2	
Blunt Force	1	1	0	0	1	3	1	0	4	2	1	0	0		14
<i>Male</i>	0	0	0	0	1	2	1	0	4	1	1	0	0	10	
<i>Female</i>	1	1	0	0	0	1	0	0	0	1	0	0	0	4	
Firearms	0	0	0	0	7	19	10	3	4	1	0	0	0		44
<i>Male</i>	0	0	0	0	4	19	10	3	4	1	0	0	0	41	
<i>Female</i>	0	0	0	0	3	0	0	0	0	0	0	0	0	3	
Homicidal Violence	0	0	0	0	0	0	0	0	0	0	0	0	0		0
<i>Male</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Stabbing	0	0	0	0	0	4	3	2	0	2	0	0	0		11
<i>Male</i>	0	0	0	0	0	3	1	2	0	2	0	0	0	8	
<i>Female</i>	0	0	0	0	0	1	2	0	0	0	0	0	0	3	
Other / Unknown	1	0	0	0	0	0	0	0	1	0	0	0	0		2
<i>Male</i>	1	0	0	0	0	0	0	0	0	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	0	0	1	0	0	0	0	1	
Totals	2	1	0	0	8	26	14	6	9	5	1	2	0		74
Percent	3%	1%	0%	0%	11%	35%	19%	8%	12%	7%	1%	3%	0%		100%

Graph 4-2 Homicide Deaths / Race / KCME / 2013



Graph 4-3 Homicide Deaths / Age Group / KCME / 2013




Table 4-3 Homicide Deaths / Age / Race / Gender / KCME / 2013

METHOD		< 16	16 to 19	20 to 29	30 to 39	40 to 49	50+	SUB-TOTAL	TOTAL
Asphyxia	White	0	0	0	0	0	2		2
	<i>Male</i>	0	0	0	0	0	1	1	
	<i>Female</i>	0	0	0	0	0	1	1	
	Asian	0	0	0	0	1	0		1
	<i>Male</i>	0	0	0	0	0	0	0	
	<i>Female</i>	0	0	0	0	1	0	1	
Blunt Force	White	0	0	2	1	0	5		8
	<i>Male</i>	0	0	2	1	0	4	7	
	<i>Female</i>	0	0	0	0	0	1	1	
	African Am.	0	0	1	0	0	2		3
	<i>Male</i>	0	0	0	0	0	2	2	
	<i>Female</i>	0	0	1	0	0	0	1	
Am. Indian / AK Native	2	1	0	0	0	0		3	
<i>Male</i>	0	1	0	0	0	0	1		
<i>Female</i>	2	0	0	0	0	0	2		
Firearms	White	0	1	10	2	2	2		17
	<i>Male</i>	0	1	9	2	2	2	16	
	<i>Female</i>	0	0	1	0	0	0	1	
	African Am.	0	3	9	6	1	2		21
	<i>Male</i>	0	3	8	6	1	2	20	
	<i>Female</i>	0	0	1	0	0	0	1	
	Asian/Pac Is.	0	0	1	2	0	0		3
	<i>Male</i>	0	0	1	2	0	0	3	
	<i>Female</i>	0	0	0	0	0	0	0	
	Am. Indian / AK Native	0	0	1	0	0	1		2
	<i>Male</i>	0	0	1	0	0	1	2	
	<i>Female</i>	0	0	0	0	0	0	0	
Other	0	0	1	0	0	0		1	
<i>Male</i>	0	0	0	0	0	0	0		
<i>Female</i>	0	0	1	0	0	0	1		
Stabbing	White	0	0	0	2	2	2		6
	<i>Male</i>	0	0	0	0	2	2	4	
	<i>Female</i>	0	0	0	2	0	0	2	
	Black	0	0	3	1	0	0		4
	<i>Male</i>	0	0	2	1	0	0	3	
	<i>Female</i>	0	0	1	0	0	0	1	
	Asian/Pac. Is	0	0	1	0	0	0		1
	<i>Male</i>	0	0	1	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	0		

Other	White	1	0	0	0	0	1	2
	Male	1	0	0	0	0	0	
	Female	0	0	0	0	0	1	
Totals		3	5	29	14	6	17	74

Table 4-4 Homicide Methods / Gender / KCME / 2013

Gender

METHOD	MALE	FEMALE	TOTAL
Asphyxia / Strangulation	1	2	3
Blunt Force	10	4	14
Firearms	41	3	44
Stabbing	8	3	11
Other / Unknown	1	1	2
Totals	61	13	74
Percent	82%	18%	100%

Table 4-5 Homicide Methods / Blood Alcohol Results / KCME / 2013

TESTED

METHOD	TESTED		NOT TESTED	TOTAL
	POSITIVE	NEGATIVE		
Asphyxia / Strangulation	2	1	0	3
Blunt Force	3	7	4	14
Firearms	12	30	2	44
Stabbing	4	6	1	11
Other / Unknown	0	0	2	2
Totals	21	44	9	74
Percent	28%	60%	12%	100%

Manner of death: Natural

The Medical Examiner assumes jurisdiction over deaths that are determined to be natural due to the sudden and unexpected nature of the death in an apparently healthy individual, when there is no physician who has knowledge or awareness of the decedent's condition, when there is no next of kin to make disposition, or when there are suspicious circumstances surrounding the death. In these situations, the Medical Examiner becomes responsible for certification of death. It should be stressed that the natural deaths the Medical Examiner investigates may not be representative of all natural deaths in the general population, due to the possibility that jurisdictional considerations introduce significant bias.

In 2013, the King County Medical Examiner's Office assumed jurisdiction over 922 deaths attributed to natural causes, representing 42% (922/2,205) of the cases investigated. The King County Medical Examiner certified 72% (661/922) of these deaths; attending physicians who had knowledge of the decedent's medical condition certified 28% (261/922). It should be noted that when a death is initially reported, there may be no evidence of an attending physician. A thorough scene investigation often reveals that the deceased did, in fact, have a physician with knowledge of the decedent's medical condition. In that case, this physician would then be contacted to certify the death.

The King County Medical Examiner performed autopsies in 72% (477/661) of the deaths certified as natural, which included autopsies performed in 100% (7/7) of deaths classified as Sudden Infant Death Syndrome (SIDS). In this context, it is important to recognize that there are changes occurring in the classification of sudden infant deaths. The term "Sudden Unexplained Infant Death" (SUID) is used by some as an alternative to SIDS. Whatever the designation, it is important to recognize that an autopsy is performed on all sudden infant deaths.

Cardiovascular disease accounted for the greatest proportion of natural deaths. Most deaths in which an autopsy was not performed were certified as due to "probable arteriosclerotic cardiovascular disease."

A special subset of deaths designated "Complication of Therapy" has been incorporated in the statistical analyses of natural deaths. Complication of Therapy is defined as a death that is identified as a predictable consequence of appropriate medical therapy. Previously, these deaths were classified separately and included in the Accident chapter. Complication of Therapy is not an official manner of death recognized by state or federal standards of death certification. It is, however, a useful category that includes deaths resulting from medical therapy or surgical procedures that are not easily classified as either natural or accidental deaths. As such, this category of deaths warrants special mention because of an apparent upward trend in incidence and increased public interest. Deaths that are excluded from this category include falls and mechanical injuries in hospitals, inadvertent misadministration of drugs, wrong-sided surgeries, and wholly unexpected procedure-related injuries, all of which are more appropriately classified as manner Accident.

As an example, a patient who dies from an infection after a colectomy for the treatment of colon cancer, that patient's death would be classified as Complication of Therapy, manner Natural. Contrast this example with the case of a patient where a proper prescription for a heart medication is written, but who is given an unintentional overdose of the medication. In this second case, the manner of death would be Accident, not Complication of Therapy.

It is important to note that the classification of a death as a Complication of Therapy is a non-judgmental means by which the inherent risk of medical therapies can be recognized and tracked. By no means is Complication of Therapy synonymous with malpractice or negligence.

Complication of Therapy can be divided into three general categories: drug-related, consequence of medical procedure, and consequence of surgery. Drug-related includes anaphylactic/allergic reaction, hemorrhagic complications of anticoagulants, anesthesia related events, and other adverse drug reactions. Consequence of medical procedure refers to complications from procedures that are therapeutic or diagnostic, but do not meet the criteria for surgery, such as placement of catheters, penetration of body cavities by needles, or manipulation of body regions, etc. Consequence of surgery refers to direct anatomic damage during a procedure and usually involves a diseased organ system, such as perforation of a viscus or vessel or hemorrhagic complications of surgery.

Graph 5-4 shows the Complication of Therapy deaths by general category and Graph 5-5 further divides the general category of surgical injury into "type of surgery" and "comorbidity." (Comorbidity is defined as the coexistence of natural disease serious enough to be listed on the death certificate as a contributing condition.)

Recognition of the importance of identifying and reporting these deaths by the medical community has surged since the Institute of Medicine of the National Academy of Sciences published a report in 1999 that estimated that up to 98,000 preventable deaths may occur each year in the United States due to medical errors. The subsequent public interest and efforts by the healthcare system to address issues of patient safety may contribute to a greater percentage of these cases being reported to the Medical Examiner.

Table 5-1 Disease Processes Causing Natural Deaths / KCME / 2013

NUMBER OF DEATHS	DISEASE DESCRIPTION
CARDIOVASCULAR	
3	Aortic aneurysm
5	Aortic dissection
87	Arteriosclerotic cardiovascular disease (ASCVD)
4	Bacterial endocarditis
7	Cardiac dysrhythmia
37	Cardiomyopathy
3	Congenital heart disease
4	Congestive heart failure
177	Hypertensive ASCVD / Hypertensive heart disease
1	Myocarditis
144	Probable arteriosclerotic cardiovascular disease
8	Valvular heart disease
3	Other
483	TOTAL CARDIOVASCULAR
CENTRAL NERVOUS SYSTEM	
18	Epilepsy (idiopathic & other non-traumatic etiologies)
7	Infarct
3	Meningitis
10	Spontaneous intracerebral hemorrhage
6	Spontaneous rupture of aneurysm
27	Other
71	TOTAL CENTRAL NERVOUS SYSTEM
COMPLICATION OF THERAPY (COT)	
5	Drug Related COT
6	Procedure Related COT
7	Surgery Related COT
18	TOTAL COMPLICATION OF THERAPY
ENDOCRINE	
5	Diabetic ketoacidosis
13	Diabetes mellitus
0	Disease of blood or blood-forming organ
0	Pancreatitis
3	Other
21	TOTAL ENDOCRINE

Table 5-1 Disease Processes Causing Natural Deaths / KCME / 2013

NUMBER OF DEATHS	DISEASE DESCRIPTION
GASTROINTESTINAL	
7	Gastrointestinal hemorrhage
3	Obstruction
1	Perforating ulcer
6	Other
17	TOTAL GASTROINTESTINAL
HEPATIC	
17	Cirrhosis
11	Cirrhosis and fatty liver
6	Fatty liver
3	Hepatic failure
15	Hepatitis
1	Other
53	TOTAL HEPATIC
MALIGNANCY	
3	Breast
2	Colon
23	Lung
4	Pancreas
4	Prostate
1	Rectum
1	Unknown
30	Other
68	TOTAL MALIGNANCY
RESPIRATORY	
0	Asthma
22	Chronic obstructive pulmonary disease
36	Pneumonia
22	Pulmonary thromboembolus
5	Other
85	TOTAL RESPIRATORY
SUDDEN INFANT DEATH SYNDROME	
7	SIDS

Table 5-1 Disease Processes Causing Natural Deaths / KCME / 2013 (continued)

NUMBER OF DEATHS	DISEASE DESCRIPTION
OTHER PROCESSES	
36	Chronic ethanolism (alcoholism)
1	Chronic prescription drug abuse
2	HIV / AIDS
1	Infant death
13	Infection
1	Intra-uterine growth retardation
7	No anatomic or toxicological cause of death
15	Sepsis
1	SUDC
22	Other
99	TOTAL OTHER PROCESSES
439	TOTAL Non-Cardiovascular Cause of Death
483	TOTAL Cardiovascular Cause of Death

922 Total NATURAL DEATHS under KCMEO Jurisdiction, 2013

Graph 5-1 Deaths due to Natural Causes / KCME / 2013





Table 5-2 Natural Deaths / Race / Gender / KCME / 2013

DISEASE PROCESS / GENDER	RACE					SUB-TOTAL	TOTAL
	WHITE	AFRIC AMER	ASIAN/ PAC IS	AM INDIAN/ /AK NATIVE	OTHER		
Cardiovascular	397	44	37	5	0		483
<i>Male</i>	271	31	24	5	0	331	
<i>Female</i>	126	13	13	0	0	152	
Central Nervous	57	7	6	1	0		71
<i>Male</i>	38	4	4	1	0	47	
<i>Female</i>	19	3	2	0	0	24	
Complication of Therapy	14	2	1	1	0		18
<i>Male</i>	4	2	1	0	0	7	
<i>Female</i>	10	0	0	1	0	11	
Endocrine	18	2	1	0	0		21
<i>Male</i>	12	2	1	0	0	15	
<i>Female</i>	6	0	0	0	0	6	
Gastrointestinal	14	1	2	0	0		17
<i>Male</i>	8	1	2	0	0	11	
<i>Female</i>	6	0	0	0	0	6	
Hepatic	49	1	0	3	0		53
<i>Male</i>	37	1	0	2	0	40	
<i>Female</i>	12	0	0	1	0	13	
Malignancy	57	7	3	1	0		68
<i>Male</i>	37	5	1	1	0	44	
<i>Female</i>	20	2	2	0	0	24	
Respiratory	72	10	3	0	0		85
<i>Male</i>	55	8	1	0	0	64	
<i>Female</i>	17	2	2	0	0	21	
SIDS	6	1	0	0	0		7
<i>Male</i>	5	0	0	0	0	5	
<i>Female</i>	1	1	0	0	0	2	
Other	84	9	3	3	0		99
<i>Male</i>	52	7	3	1	0	63	
<i>Female</i>	32	2	0	2	0	36	
Totals	768	84	56	14	0		922
Percent	83%	9%	6%	2%	0%		100

Graph 5-2 Natural Deaths / Race / KCME / 2013



Graph 5-3 Natural Deaths / Age Group / KCME / 2013






Table 5-3 Natural Deaths / Age / Gender / KCME / 2013

DISEASE PROCESS/ GENDER	AGE GROUP (YEARS)													SUB-TOTAL	TOTAL
	< 1	1 to 5	6 to 12	13 to 15	16 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 to 69	70 to 79	80 to 89	90 +		
Cardiovascular	1	0	0	0	0	6	13	51	99	154	83	53	23		483
<i>Male</i>	1	0	0	0	0	5	10	43	81	112	52	21	6	331	
<i>Female</i>	0	0	0	0	0	1	3	8	18	42	31	32	17	152	
Central Nervous	0	0	1	0	2	6	6	6	11	8	11	11	9		71
<i>Male</i>	0	0	0	0	2	4	3	4	9	8	7	7	3	47	
<i>Female</i>	0	0	1	0	0	2	3	2	2	0	4	4	6	24	
Complication of Therapy	0	2	0	0	0	1	2	1	1	5	4	1	1		18
<i>Male</i>	0	2	0	0	0	0	1	0	0	1	2	1	0	7	
<i>Female</i>	0	0	0	0	0	1	1	1	1	4	2	0	1	11	
Endocrine	0	0	0	0	0	0	1	4	7	7	2	0	0		21
<i>Male</i>	0	0	0	0	0	0	0	3	4	6	2	0	0	15	
<i>Female</i>	0	0	0	0	0	0	1	1	3	1	0	0	0	6	
Gastrointestinal	0	0	0	0	0	0	1	3	6	6	0	1	0		17
<i>Male</i>	0	0	0	0	0	0	1	2	4	3	0	1	0	11	
<i>Female</i>	0	0	0	0	0	0	0	1	2	3	0	0	0	6	
Hepatic	0	0	0	0	0	1	5	11	24	10	2	0	0		53
<i>Male</i>	0	0	0	0	0	1	1	8	22	7	1	0	0	40	
<i>Female</i>	0	0	0	0	0	0	4	3	2	3	1	0	0	13	
Malignancy	0	0	0	0	0	0	1	2	17	20	21	5	2		68
<i>Male</i>	0	0	0	0	0	0	0	2	12	12	13	4	1	44	
<i>Female</i>	0	0	0	0	0	0	1	0	5	8	8	1	1	24	
Respiratory	3	2	0	0	0	0	3	10	19	20	17	9	2		85
<i>Male</i>	3	1	0	0	0	0	2	7	16	16	12	6	1	64	
<i>Female</i>	0	1	0	0	0	0	1	3	3	4	5	3	1	21	
SIDS	7	0	0	0	0	0	0	0	0	0	0	0	0		7
<i>Male</i>	5	0	0	0	0	0	0	0	0	0	0	0	0	5	
<i>Female</i>	2	0	0	0	0	0	0	0	0	0	0	0	0	2	
Other	2	3	0	0	1	5	9	16	26	23	10	3	1		99
<i>Male</i>	0	3	0	0	1	2	5	9	17	16	8	2	0	63	
<i>Female</i>	2	0	0	0	0	3	4	7	9	7	2	1	1	36	
Totals	13	7	1	0	3	19	41	104	210	253	150	83	38		922
Percent	1.4%	0.8%	0.1%	0%	0.3%	2%	4.5%	11.3%	22.8%	27.4%	16.3%	9%	4.1%		100%




Table 5-4 Natural Deaths / Gender / KCME / 2013

CIRCUMSTANCES	GENDER		TOTAL
	MALE	FEMALE	
Cardiovascular	331	152	483
Central Nervous	47	24	71
Complication of Therapy	7	11	18
Endocrine	15	6	21
Gastrointestinal	11	6	17
Hepatic	40	13	53
Malignancy	44	24	68
Respiratory	64	21	85
SIDS	5	2	7
Other	63	36	99
Totals	627	295	922
Percent	68%	32%	100%


Table 5-5 Natural Deaths / Blood Alcohol Results / KCME / 2013

METHOD	TESTED		NOT	TOTAL
	POSITIVE	NEGATIVE	TESTED	
Cardiovascular	56	238	189	483
Central Nervous System	2	30	39	71
Complication of Therapy	1	8	9	18
Endocrine	1	7	13	21
Gastrointestinal	6	4	7	17
Hepatic	19	19	15	53
Malignancy	1	10	57	68
Respiratory	6	43	36	85
SIDS	0	7	0	7
Other	18	24	57	99
Totals	110	390	422	922
Percent	12%	42%	46%	100%

Graph 5-4 Complication of Therapy / General Categories / KCME / 2013



Graph 5-5 Complication of Therapy¹⁶ / Surgical Injuries / KCME / 2013



¹⁶Serious co-morbidity indicates coexisting natural disease serious enough to contribute to death.

Manner of death: Suicide

Suicides are deaths caused by self-inflicted injuries with evidence of intent to end one's life. Evidence of intent includes an explicit expression, such as a suicide note or verbal threat, or an act constituting implicit intent, such as deliberately placing a gun to one's head or rigging a vehicle's exhaust. In 2013, there were 266 suicides, accounting for 12% (266/2,205) of the deaths that the King County Medical Examiner's Office investigated.

In 2013, there were fifteen suicides among persons 19 years and younger (5.6% of all suicides, 15/266), which is lower than in 2012 when there were eighteen suicides in this age group. Suicides in the age group 60 years and older represented 32% (84/266) of all suicides in 2013.

Firearms were responsible for 38% (100/266) of the 2013 suicide deaths. The number of gunshot suicides (100/266) in 2013 is nineteen less than in 2012 when there were 119. Hanging accounted for 27% (71/266) of suicidal deaths, while jumping from a height accounted for 5.6% (15/266). Drugs and poisons accounted for 15% (41/266) of all suicides, while carbon monoxide caused death in 4% (10/266) of the cases. More information regarding drug-caused deaths is presented in the section "Deaths Due to Drugs & Poisons" beginning on page 89.

Firearms were the primary method of committing suicide for all age groups over the age of 20. In the 19 years and younger age group, firearms represented 6.7% (1/15) of the deaths while hanging represented 73% (11/15) of the deaths.

Blood alcohol tests were performed in 97% (257/266) of suicidal deaths and were positive in 34% (88/257) of cases tested.

In 2013, there were eleven deaths due to drugs and/or poisons by adults 60 years of age and over. In 2013, there was one suicide attributed to drugs and/or poisons among youths 19 years and younger. In 2012, there were no deaths from drug and/or poisons in this age group.

The [Washington Death with Dignity Act, Initiative 1000](#), codified as [RCW 70.245](#), passed on November 4, 2008 and took effect on March 5, 2013. This act allows terminally ill adults seeking to end their life to request lethal doses of medication from medical and osteopathic physicians. These terminally ill patients must be Washington state residents who have less than six months to live.¹⁷

As provided in the act, "the patient's death certificate...shall list the underlying terminal disease as the cause of death." The act also states that, "Actions taken in accordance with this chapter do not, for any purpose, constitute suicide, assisted suicide, mercy killing, or homicide, under the law." Given these instructions, the King County Medical Examiner's Office has no involvement in these cases and collects no statistics on the number of deaths where an individual has utilized their rights under the provisions of this act. Statistics are kept and released annually by the Washington State Department of Health.

¹⁷ Washington State Department of Health website: <http://www.doh.wa.gov/dwda>

Graph 6-1 Suicide Injury Methods / KCME / 2013







Table 6-1 Suicide Injury Methods / Race / Gender / KCME / 2013

CIRCUMSTANCES / GENDER	RACE					SUB-TOTAL	TOTAL
	WHITE	AFRIC AMER	ASIAN/ PAC IS	AM INDIAN/ AK NATIVE	OTHER		
Asphyxia	12	1	0	0	0		13
<i>Male</i>	7	1	0	0	0	8	
<i>Female</i>	5	0	0	0	0	5	
Burns / Fire	1	0	0	0	0		1
<i>Male</i>	1	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	
Carbon Monoxide	8	2	0	0	0		10
<i>Male</i>	8	1	0	0	0	9	
<i>Female</i>	0	1	0	0	0	1	
Drowning	1	0	1	0	0		2
<i>Male</i>	0	0	1	0	0	1	
<i>Female</i>	1	0	0	0	0	1	
Drugs / Poisons	38	0	3	0	0		41
<i>Male</i>	18	0	0	0	0	18	
<i>Female</i>	20	0	3	0	0	23	
Firearms	85	6	8	1	0		100
<i>Male</i>	74	4	8	1	0	87	
<i>Female</i>	11	2	0	0	0	13	
Hanging	55	1	13	1	1		71
<i>Male</i>	40	1	10	1	1	53	
<i>Female</i>	15	0	3	0	0	18	
Incised / Stab Wound(s)	8	0	1	0	0		9
<i>Male</i>	7	0	0	0	0	7	
<i>Female</i>	1	0	1	0	0	2	
Jumping	14	1	0	0	0		15
<i>Male</i>	9	1	0	0	0	10	
<i>Female</i>	5	0	0	0	0	5	
Other	1	0	0	0	0		1
<i>Male</i>	0	0	0	0	0	0	
<i>Female</i>	1	0	0	0	0	1	
Traffic	2	0	0	1	0		3
<i>Male</i>	2	0	0	1	0	3	
<i>Female</i>	0	0	0	0	0	0	
Totals	225	11	26	3	1		266
Percent	84.6%	4%	10%	1%	0.4%		100%

Graph 6-2 Suicide Deaths / Race / KCME / 2013



Graph 6-3 Suicide Deaths / Age Group / KCME / 2013




Table 6-2

Suicide Injury Methods / Age / Gender / KCME / 2013

King County Medical Examiner's Office - 2013 Annual Report

INJURY METHOD/ GENDER	AGE GROUP (YEARS)									SUB-TOTAL	TOTAL
	12 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 to 69	70 to 79	80 to 89	90 +		
Asphyxia	0	1	2	2	2	0	2	3	1		13
<i>Male</i>	0	1	2	1	0	0	1	2	1	8	
<i>Female</i>	0	0	0	1	2	0	1	1	0	5	
Burns / Fire	0	0	0	0	1	0	0	0	0		1
<i>Male</i>	0	0	0	0	1	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	
Carbon Monoxide	1	0	2	1	3	3	0	0	0		10
<i>Male</i>	1	0	2	0	3	3	0	0	0	9	
<i>Female</i>	0	0	0	1	0	0	0	0	0	1	
Drowning	0	1	0	0	0	0	0	1	0		2
<i>Male</i>	0	1	0	0	0	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	0	1	0	1	
Drugs / Poisons	1	2	6	13	8	9	1	0	1		41
<i>Male</i>	1	1	1	6	3	5	0	0	1	18	
<i>Female</i>	0	1	5	7	5	4	1	0	0	23	
Firearms	1	15	15	7	19	23	9	8	3		100
<i>Male</i>	1	12	13	6	16	19	9	8	3	87	
<i>Female</i>	0	3	2	1	3	4	0	0	0	13	
Hanging	11	16	15	9	8	8	3	0	1		71
<i>Male</i>	9	11	14	6	6	4	2	0	1	53	
<i>Female</i>	2	5	1	3	2	4	1	0	0	18	
Incised / Stab Wound(s)	0	1	1	1	2	2	0	2	0		9
<i>Male</i>	0	0	1	1	2	2	0	1	0	7	
<i>Female</i>	0	1	0	0	0	0	0	1	0	2	
Jumping	1	3	3	2	3	1	2	0	0		15
<i>Male</i>	1	2	3	1	2	1	0	0	0	10	
<i>Female</i>	0	1	0	1	1	0	2	0	0	5	
Other	0	0	0	0	0	1	0	0	0		1
<i>Male</i>	0	0	0	0	0	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	1	0	0	0	1	
Traffic	0	0	1	1	1	0	0	0	0		3
<i>Male</i>	0	0	1	1	1	0	0	0	0	3	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	
Totals	15	39	45	36	47	47	17	14	6		266
Percent	5.6%	14.7%	17%	13.5%	17.6%	17.6%	6.4%	5.3%	2.3%		100%

Table 6-3

Suicide Injury Methods / Gender / KCME / 2013



INJURY METHOD	GENDER		TOTAL
	MALE	FEMALE	
Asphyxia	8	5	13
Burns/ Fire	1	0	1
Carbon Monoxide	9	1	10
Drowning	1	1	2
Drugs / Poisons	18	23	41
Firearms	87	13	100
Hanging	53	18	71
Incised / Stab Wound(s)	7	2	9
Jumping	10	5	15
Other	0	1	1
Traffic	3	0	3
Totals	197	69	266
Percent	74%	26%	100%

Table 6-4

Suicide Injury Methods / Marital Status / Gender / KCME / 2013

King County Medical Examiner's Office - 2013 Annual Report

CIRCUMSTANCES / GENDER	MARITAL STATUS					Sub-Total	Total
	Never Married	Married	Divorced	Widowed	Unknown		
Asphyxia	5	6	2	0	0		13
<i>Male</i>	3	4	1	0	0	8	
<i>Female</i>	2	2	1	0	0	5	
Burns/ Fire	0	1	0	0	0		1
<i>Male</i>	0	1	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	
Carbon Monoxide	4	3	2	0	1		10
<i>Male</i>	3	3	2	0	1	9	
<i>Female</i>	1	0	0	0	0	1	
Drowning	1	0	1	0	0		2
<i>Male</i>	1	0	0	0	0	1	
<i>Female</i>	0	0	1	0	0	1	
Drugs / Poisons	18	8	14	1	0		41
<i>Male</i>	10	3	4	1	0	18	
<i>Female</i>	8	5	10	0	0	23	
Firearms	36	38	21	5	0		100
<i>Male</i>	32	32	19	4	0	87	
<i>Female</i>	4	6	2	1	0	13	
Hanging	42	15	10	2	2		71
<i>Male</i>	33	10	8	0	2	53	
<i>Female</i>	9	5	2	2	0	18	
Incised / Stab Wound(s)	3	3	3	0	0		9
<i>Male</i>	2	3	2	0	0	7	
<i>Female</i>	1	0	1	0	0	2	
Jumping	8	4	3	0	0		15
<i>Male</i>	5	3	2	0	0	10	
<i>Female</i>	3	1	1	0	0	5	
Other	0	0	1	0	0		1
<i>Male</i>	0	0	0	0	0	0	
<i>Female</i>	0	0	1	0	0	1	
Traffic	1	0	2	0	0		3
<i>Male</i>	1	0	2	0	0	3	
<i>Female</i>	0	0	0	0	0	0	
Totals	118	78	59	8	3		266
Percent	44.4%	29.3%	22.2%	3.0%	1.1%		100%

Table 6-5

Suicide Injury Methods / Blood Alcohol / KCME / 2013

King County Medical Examiner's Office - 2013 Annual Report



METHOD	TESTED		NOT TESTED	TOTAL
	POSITIVE	NEGATIVE		
Asphyxia	4	9	0	13
Burns/ Fire	0	0	1	1
Carbon Monoxide	6	3	1	10
Drowning	0	2	0	2
Drugs / Poisons	15	26	0	41
Firearms	37	59	4	100
Hanging	22	47	2	71
Incised / Stab Wound(s)	1	8	0	9
Jumping	2	12	1	15
Other	0	1	0	1
Traffic	1	2	0	3
Totals	88	169	9	266
Percent	33%	64%	3%	100%


Manner of death: Undetermined

The King County Medical Examiner's Office certifies a manner of death as undetermined when available information regarding the circumstances of death is insufficient to classify the death into one of the specific manners of natural or unnatural (Accident, Homicide or Suicide) death. In some cases, serious doubt exists as to whether an injury occurred with intent or as a result of an accident. Information concerning the circumstances may be lacking due to the absence of background information or witnesses, or because of a lengthy delay between death and discovery of the body. Moreover, it may be difficult to assess street drug or medication overdose deaths as showing enough features to reasonably determine the manner of death. If an extensive investigation and autopsy cannot clarify the circumstances, the death is classified undetermined.

The King County Medical Examiner's Office certified 66 deaths with manner undetermined, accounting for 3% (66/2,119) of the deaths investigated in 2013. Drugs and poisons caused 11% (7/66) of the deaths classified as undetermined. For a more detailed review of drug-caused deaths in 2013, see the discussion in the section on Drugs and Poisons on pages 89 and 90.

The 66 deaths that were classified as undetermined for 2013 included 5 fetal deaths, which, in accordance with the Washington State Department of Health - Center for Health Statistics Fetal Death Certification Guidelines, are not assigned a manner of death. Fetal death certificates must be issued for every fetus of 20 weeks or more gestation. Of the 5 fetal deaths in 2013, one was related to maternal drug abuse.

Graph 7-1 Undetermined Manner of Death¹⁸ / KCME / 2013




¹⁸NATCOD is an abbreviation for “no anatomic or toxicological cause of death,” and refers to deaths in which full autopsies and toxicological analyses (if relevant) fail to identify an adequate cause of death.



Table 7-1 Undetermined Manner of Death / Race / Gender / KCME / 2013

CIRCUMSTANCES / GENDER	RACE					SUB-TOTAL	TOTAL
	WHITE	AFRIC AMER	ASIAN/ PAC IS	AM INDIAN/ AK NATIVE	OTHER / UNK		
Blunt Force	6	2	0	2	0		10
<i>Male</i>	4	2	0	1	0	7	
<i>Female</i>	2	0	0	1	0	3	
Burns / Fire	0	0	0	0	0		1
<i>Male</i>	0	0	0	1	0	1	
<i>Female</i>	0	0	0	0	0	0	
Drugs / Poisons	5	1	1	0	0		7
<i>Male</i>	4	1	0	0	0	5	
<i>Female</i>	1	0	1	0	0	2	
Fall	4	1	1	0	0		6
<i>Male</i>	3	1	1	0	0	5	
<i>Female</i>	1	0	0	0	0	1	
Fetal Deaths	3	0	0	2	0		5
<i>Male</i>	2	0	0	1	0	3	
<i>Female</i>	1	0	0	1	0	2	
Firearms	0	0	0	0	0		0
<i>Male</i>	0	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	0	
No Anatomic or Toxicological Cause of Death	4	1	1	0	0		6
<i>Male</i>	2	0	1	0	0	3	
<i>Female</i>	2	1	0	0	0	3	
Other	13	2	2	3	0		20
<i>Male</i>	9	2	2	2	0	15	
<i>Female</i>	4	0	0	1	0	5	
SIDS	7	2	1	1	0		11
<i>Male</i>	4	1	1	1	0	7	
<i>Female</i>	3	1	0	0	0	4	
Totals	42	9	6	9	0		11
Percent	64%	13.5%	9%	13.5%	0%		100%

Graph 7-2 Undetermined Manner / Race / KCME / 2013



Graph 7-3 Undetermined Manner / Age Group / KCME / 2013






Table 7-2 Undetermined Circumstances / Age / Gender / KCME / 2013

INJURY METHOD / GENDER	AGE GROUP (YEARS)												SUB-TOTAL	TOTAL
	<1	1 to 5	6 to 15	16 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 to 69	70 to 79	80 to 89	90 +		
Blunt Force	1	1	0	1	0	1	2	4	0	0	0	0		10
<i>Male</i>	1	1	0	1	0	0	1	4	0	0	0	0	8	
<i>Female</i>	0	0	0	0	0	1	1	0	0	0	0	0	2	
Burns / Fire	0	0	0	0	0	0	0	1	0	0	0	0		1
<i>Male</i>	0	0	0	0	0	0	0	1	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	
Drugs / Poisons	0	0	0	0	0	4	3	0	0	0	0	0		7
<i>Male</i>	0	0	0	0	0	3	2	0	0	0	0	0	5	
<i>Female</i>	0	0	0	0	0	1	1	0	0	0	0	0	2	
Fall	0	0	0	0	1	1	2	1	0	0	0	1		6
<i>Male</i>	0	0	0	0	1	1	2	1	0	0	0	0	5	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	1	1	
Fetal Deaths	5	0	0	0	0	0	0	0	0	0	0	0		5
<i>Male</i>	3	0	0	0	0	0	0	0	0	0	0	0	3	
<i>Female</i>	2	0	0	0	0	0	0	0	0	0	0	0	2	
Firearms	0	0	0	0	0	0	0	0	0	0	0	0		0
<i>Male</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	
No anatomic or toxicological cause of death	0	0	0	0	1	1	2	1	1	0	0	0		6
<i>Male</i>	0	0	0	0	1	0	2	0	0	0	0	0	3	
<i>Female</i>	0	0	0	0	0	1	0	1	1	0	0	0	3	
Other	0	0	0	0	3	1	5	3	1	0	5	2		20
<i>Male</i>	0	0	0	0	3	1	4	3	0	0	2	2	15	
<i>Female</i>	0	0	0	0	0	0	1	0	1	0	3	0	5	
SIDS	11	0	0	0	0	0	0	0	0	0	0	0		11
<i>Male</i>	7	0	0	0	0	0	0	0	0	0	0	0	7	
<i>Female</i>	4	0	0	0	0	0	0	0	0	0	0	0	4	
Totals	17	1	0	1	5	8	14	10	2	0	5	3		66
Percent	26%	1%	0%	1%	8%	12%	21%	15%	3%	0%	8%	5%		100%

Table 7-3 Undetermined Manner / Gender / KCME / 2013



INJURY METHOD	GENDER		TOTAL
	MALE	FEMALE	
Blunt Force	7	3	10
Burns / Fire	1	0	1
Drugs / Poisons	5	2	7
Fall	5	1	6
Fetal Deaths	3	2	5
Firearms	0	0	0
No Anatomic or Toxicological Cause of Death	3	3	6
Other	15	5	20
SIDS	7	4	11
Totals	46	20	66
Percent	70%	30%	100%

Table 7-4 Undetermined Manner / Blood Alcohol Results / KCME / 2013

METHOD	TESTED		NOT TESTED	TOTAL
	POSITIVE	NEGATIVE		
Blunt Force	4	5	1	10
Burns / Fire	1	0	0	1
Drugs / Poisons	1	6	0	7
Fall	2	3	1	6
Fetal Deaths	0	2	3	5
Firearms	0	0	0	0
No Anatomic or Toxicological Cause of Death	1	3	2	6
Other	6	11	3	20
SIDS	0	9	2	11
Totals	15	39	12	66
Percent	23%	59%	18%	100%

Traffic deaths

During the calendar year 2013, the Medical Examiner's Office participated in the investigation of 123 traffic fatalities. In 67% (82/123) of the traffic deaths, the collisions occurred in King County, compared to 74% (97/131) of the collisions in 2012. In 2013, 33% (41/123) of the traffic deaths that the Medical Examiner investigated were the result of collisions that occurred outside of King County, with the injured transported to hospitals in King County, primarily Harborview Medical Center. Because the deaths occurred in King County, it falls under the jurisdiction of the King County Medical Examiner. Although these deaths are classified "Accident" for death certification purposes, the more accurate term is "motor vehicle collision."

In 2013, 37% (45/123) of the traffic fatalities were motor vehicle drivers. Teenage drivers (16-19 years of age) were 20% (9/45) of the driver deaths in 2013 and 9% (4/47) in 2012. By age, 56% of vehicle driver deaths (25/45) were people between the ages of 20 and 29. Thirteen percent of driver deaths (6/45) were adults between the ages of 30 and 39. Thirty-three percent (15/45) were adults between the ages of 40 and 49. Male drivers represented 69% (31/45) of driver deaths and female drivers represented 31% of driver deaths (14/45).

Of the 123 traffic fatalities in 2013, 23 were motor vehicle passengers, representing 19% of the total (23/123). In 2013, teenagers (13-19 years old) accounted for 2 motor vehicle passenger deaths. There were no passenger deaths of infants (less than one year of age), 1 vehicle passenger death of a child between the ages of 1-5 years, and no deaths of children between the ages of 6-12 years.

Blood ethanol (alcohol) statistics are presented to describe the role of alcohol in traffic deaths. However, it should be noted that in many cases someone other than the person who died was under the influence of alcohol and was directly responsible for the collision. The Medical Examiner determines the blood alcohol levels of persons who die, not of everyone involved in the incident. In addition, blood alcohol is not tested in persons who die after surviving more than 24 hours, because in those deaths the alcohol has had time to metabolize.¹⁹ Therefore, blood alcohol figures presented in this report are not a total description of the role of alcohol in traffic collisions.

Of cases in which seatbelt restraint status was known, 20% (9/45) of drivers in vehicle deaths were not restrained. The figures for drivers not wearing seatbelts for the previous three years are: 26% (10/38) in 2013, 33% (13/39) in 2012, and 35% (23/65) in 2011.

Motorcycle riders accounted for 18% (22/123) of traffic fatalities. In 2013, there were 22 motorcycle driver fatalities and no motorcycle passenger fatalities. All twenty-two of the motorcycle driver deaths were male and none were female. Of the 22 motorcycle fatalities, 73% (16/22) of the motorcyclists were wearing a helmet; in one case, it was unknown if the motorcycle driver was wearing a helmet. Twenty of the motorcyclist fatalities were tested for the presence of blood alcohol. Six, or 30% (6/20), had a detectable amount of alcohol at the time of autopsy.

Pedestrians constituted 20% (25/123) of traffic fatalities. The majority of pedestrian deaths, 68% (17/25), were male. Of the pedestrian fatalities that were tested, 44% (7/16) had detectable amounts of alcohol present in their blood at the time of death.

¹⁹See "Explanation of Data" for criteria for blood alcohol testing, page 6.



There were seven bicyclist deaths in 2013; 3 riders were wearing helmets, 2 were not wearing a helmet, and it is unknown if 2 were wearing a helmet or not. Of the bicyclist fatalities that were tested, none had a detectable amount of alcohol present in his/her blood at the time of death.

Graph 8-1 Traffic Fatality Circumstances / KCME / 2013







Table 8-1 Traffic Fatality Circumstances / Race / Gender / KCME / 2013

CIRCUMSTANCES / GENDER	RACE					SUB-TOTAL	TOTAL
	WHITE	AFRICAN AMER	ASIAN/ PAC IS	AM INDIAN /AK NATIVE	OTHER		
Vehicle Driver	39	1	4	1	0		45
<i>Male</i>	26	1	3	1	0	31	
<i>Female</i>	13	0	1	0	0	14	
Vehicle Passenger	17	6	0	0	0		23
<i>Male</i>	7	4	0	0	0	11	
<i>Female</i>	10	2	0	0	0	12	
Vehicle Unknown Position	0	0	0	0	0		0
<i>Male</i>	0	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	0	
Bicycle	5	0	0	2	0		7
<i>Male</i>	5	0	0	2	0	7	
<i>Female</i>	0	0	0	0	0	0	
Motorcycle Driver	21	1	0	0	0		22
<i>Male</i>	21	1	0	0	0	22	
<i>Female</i>	0	0	0	0	0	0	
Motorcycle Passenger	0	0	0	0	0		0
<i>Male</i>	0	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	0	
Pedestrian	19	2	2	2	0		25
<i>Male</i>	12	2	1	2	0	17	
<i>Female</i>	7	0	1	0	0	8	
Other	1	0	0	0	0		1
<i>Male</i>	1	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	
Totals	102	10	6	5	0		123
Percent	83%	8%	5%	4%	0%		100%

Graph 8-2 Traffic Fatalities / Race / KCME / 2013



Graph 8-3 Traffic Fatalities / Age / KCME / 2013






Table 8-2 Traffic Fatality Circumstances / Age / Gender / KCME / 2013

Circumstances/Gender	AGE GROUP (YEARS)													SUB-TOTAL	TOTAL
	< 1	1 to 5	6 to 12	13 to 15	16 to 19	20 to 29	30 to 39	40 to 49	50 to 59	60 to 69	70 to 79	80 to 89	90 +		
Vehicle Driver	0	0	0	0	7	10	1	3	7	2	5	8	2		45
<i>Male</i>	0	0	0	0	6	8	1	3	4	2	3	4	1	31	
<i>Female</i>	0	0	0	0	1	2	0	1	3	0	2	4	1	14	
Vehicle Passenger	0	1	0	1	1	11	0	2	2	0	2	1	2		23
<i>Male</i>	0	1	0	0	0	5	0	2	1	0	1	0	1	11	
<i>Female</i>	0	0	0	1	1	6	0	0	1	0	1	1	1	12	
Vehicle Position Unknown	0	0	0	0	0	0	0	0	0	0	0	0	0		0
<i>Male</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Bicyclist	0	0	0	0	0	0	0	3	3	0	0	1	0		7
<i>Male</i>	0	0	0	0	0	0	0	3	3	0	0	1	0	7	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Motorcycle Driver	0	0	0	1	0	3	3	4	6	5	0	0	0		22
<i>Male</i>	0	0	0	1	0	3	3	4	6	5	0	0	0	22	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Motorcycle Passenger	0	0	0	0	0	0	0	0	0	0	0	0	0		0
<i>Male</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Pedestrian	0	0	0	1	0	1	2	3	2	8	6	2	0		25
<i>Male</i>	0	0	0	0	0	1	2	3	1	5	4	1	0	17	
<i>Female</i>	0	0	0	1	0	0	0	0	1	3	2	1	0	8	
Other	0	0	0	0	1	0	0	0	0	0	0	0	0		1
<i>Male</i>	0	0	0	0	1	0	0	0	0	0	0	0	0	1	
<i>Female</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Totals	0	1	0	3	9	25	6	15	20	15	13	12	4		123
Percent	0%	0.8%	0%	2.4%	7.3%	20.3%	4.8%	12.2%	16.3%	12.2%	10.6%	9.8%	3.3%		100%



Table 8-3 Traffic Fatality Circumstances / Gender / KCME / 2013

CIRCUMSTANCES	GENDER		TOTAL
	MALE	FEMALE	
Vehicle Driver	31	14	45
Vehicle Passenger	11	12	23
Vehicle Position Unknown	0	0	0
Bicyclist	7	0	7
Motorcycle Driver	22	0	22
Motorcycle Passenger	0	0	0
Pedestrian	17	8	25
Other Mode	1	0	1
Totals	89	34	123
Percent	72%	28%	100%

Table 8-4 Traffic Fatality Circumstances / Use of Restraint / Helmet / KCME / 2013²

CIRCUMSTANCES	Used Safety Device	No Safety Device Used	Unknown	TOTAL
Vehicle Driver	25	9	11	45
Vehicle Passenger	10	10	3	23
Bicyclist	3	2	2	7
Motorcycle Driver	16	5	1	22
Motorcycle Passenger	0	0	0	0
Totals	54	26	17	97
Percent	56%	27%	17%	100%

²Does not include vehicle position unknown, pedestrian or other traffic modes of deaths.



Table 8-5 Traffic Fatality Circumstances / Blood Alcohol Results / KCME / 2013


CIRCUMSTANCES	TESTED		NOT TESTED	TOTAL
	POSITIVE	NEGATIVE		
Vehicle Driver	7	25	13	45
Vehicle Passenger	6	12	5	23
Vehicle Position Unknown	0	0	0	0
Bicyclist	0	5	2	7
Motorcycle Driver	6	14	2	22
Motorcycle Passenger	0	0	0	0
Pedestrian	7	9	9	25
Other Mode	0	0	1	1
Totals	26	65	32	123
Percent	21%	53%	26%	100%



Table 8-6 Time of Fatal Traffic Collision / KCME / 2013

TIME OF DAY	TOTAL	PERCENT
12:01 a.m. - 3:00 a.m.	11	9%
3:01 a.m. - 6:00 a.m.	11	9%
6:01 a.m. - 9:00 a.m.	11	9%
9:01 a.m. - Noon	7	6%
12:01 p.m. - 3:00 p.m.	18	15%
3:01 p.m. - 6:00 p.m.	25	20%
6:01 p.m. - 9:00 p.m.	19	15%
9:01 p.m. - Midnight	15	12%
Unknown	6	5%
TOTALS	123	100%

Graph 8-5 Time of Fatal Traffic Collision / KCME / 2013



Deaths due to drugs and poisons

In 2012, it was reported in the *National Vital Statistics Report*²⁰ that preliminary cause of death information from 2009 shows drug-induced deaths were the leading cause of accidental deaths of Americans. This was the first time drug-induced deaths had surpassed motor vehicle accidents as the number one cause of accidental deaths.

For King County in 2013, drugs and poisons caused 329 deaths, approximately 16% of all deaths investigated (329/2,119). The total number of drug-caused deaths increased compared to 2012 when there were 298 drug deaths. In 2013, deaths due to drugs and poisons comprised 29% (329/1123) of all suicidal, accidental and undetermined deaths combined.

For the purpose of this section, the term “overdose” is used to describe a death caused by a single drug or multiple drugs in combination. Multiple drug intoxication continued to cause the majority of drug deaths in 2013. Of the drug/poison deaths in 2013, a single drug or poison caused 33% of the drug related deaths (109/329), and drugs or poisons in combination caused 67% (220/329.) Multiple drug intoxication caused 62% of the drug/poison deaths in 2012. Table 9-3 displays the specific drugs that caused death in 2013. Because of their prevalence, ethanol, cocaine (a stimulant), and opiates²¹ are identified as separate drug categories. Data on deaths involving methadone, oxycodone, and methamphetamine are also shown in detail.

Deaths due to drugs and poisons are represented in the manners of accident, suicide, and undetermined. There were no deaths classified as homicide in 2013 in which drugs or poisons were the primary cause of the death, although the victim may have been under the influence of drugs at the time of the fatal incident.

The classification of undetermined manner is used when the circumstances surrounding the drug death does not allow clarification of whether the fatal intoxication was intentional, unintentional ("recreational"), or due to another person's actions. In the year 2013, drugs and poisons caused 9 deaths of undetermined manner, compared to 17 in 2012. Of the 9 undetermined drug related deaths in 2013, 2 were fetal deaths attributed to maternal methamphetamine use.

In 2013, drugs/poisons caused 41 suicides, as compared to 51 in 2012.

²⁰ Kenneth D. Kochanek, M.A.; Jiaquan Xu, M.D.; Sherry L. Murphy, B.S.; Arialdi M. Miniño M.P.H.; and Hsiang-Ching Kung, Ph.D., Division of Vital Statistics “Deaths: Preliminary Data 2009,” National Vital Statistics Report Volume 59 Number 4 (March 2012)

²¹ When the term “opiate” is used in this section, the drug detected by analysis is a derivative of opium, usually morphine, the source of which is either pharmaceutical morphine or heroin. The term opioid refers to the general class of drugs, often called narcotics that interact with the opioid receptor. For example, oxycodone, and methadone are “opioids” but in this section are not “opiates.”

Drugs/poisons caused 279 accidental overdose deaths in 2013 compared to 230 in 2012. In 2013, accidental drug deaths comprised 42% (279/668) of all accidental deaths.

Ethanol (alcohol) is also a drug to be critically examined for its role in the circumstances surrounding death. In 2013, 28 accidental deaths were attributed to acute ethanol intoxication where ethanol was the single substance used. Fifty-four (77) people died in 2013 where ethanol, in combination with other drugs, was the cause of death. Blood alcohol (ethanol) tests were performed in 73% (875/1197) of non-natural deaths. Blood alcohol tests are only performed when death occurs within 24 hours of the initial injury/event, or, in hospital deaths, when an admission blood sample is available for testing. Positive blood alcohol levels were detected in 23% (270/1197) of non-natural deaths where tests were performed.

Table 9-1 Blood Alcohol Testing / Manner / KCME / 2013

Test Results	ACCIDENT	TRAFFIC	HOMICIDE	NATURAL	SUICIDE	UNDETERMINED	TOTAL
Tested	403	96	65	506	257	54	1381
<i>Positive</i>	119	27	21	110	88	153	380
<i>Negative</i>	284	69	44	396	169	39	1001
Not Tested	265	27	9	416	9	12	738
Totals	668	123	74	922	266	66	2119

Table 9-2 Blood Alcohol Testing / Percentage / Manner / KCME / 2013

Test Results	ACCIDENT	TRAFFIC	HOMICIDE	NATURAL	SUICIDE	UNDETERMINED	TOTAL
Tested	60%	78%	88%	55%	97%	82%	65%
<i>Positive</i>	18%	22%	28%	12%	33%	23%	18%
<i>Negative</i>	43%	56%	60%	43%	64%	54%	47%
Not Tested	40%	22%	12%	45%	3%	18%	35%
Totals	100%	100%	100%	100%	100%	100%	100%

Table 9-3 2013 Drug & Poison Caused Deaths¹

Drug Name	Total deaths out of 2,119 cases in which drug was present	Overdose Deaths (329) – Drug Present						Overdose Deaths (329) – Drug Causing					
		In which drug was present	Single drug OD in which drug was present	Multiple drug OD in which drug was present	Accident	Suicide	Undetermined	In which drug caused death	OD in which a single drug caused death	OD in which multiple drugs caused death	Accident	Suicide	Undetermined
Acetaminophen	6	2	1	1	1	1	0	1	1	0	1	0	0
Alprazolam	38	27	0	27	25	2	0	26	0	26	24	2	0
Amitriptyline	17	10	1	9	5	5	0	10	1	9	5	5	0
Amphetamine	124	36	12	24	22	0	2	35	12	23	21	0	2
Bupropion	8	4	1	3	2	1	1	3	0	3	1	1	1
Butalbital	7	3	0	3	2	1	0	3	0	3	2	1	0
Cannabinoids / THC ²	105	23	5	18	22	1	0	0	0	0	0	0	0
Carbamazepine	7	2	0	2	1	0	1	1	0	1	0	0	1
Carbon Monoxide ³	28	2	1	1	1	1	0	0	0	0	0	0	0
Carisoprodol	4	4	0	4	4	0	0	4	0	4	4	0	0
Chlordiazepoxide	13	7	1	6	6	0	1	6	0	6	5	0	1
Chlorpromazine	1	1	0	1	1	0	0	1	0	1	1	0	0
Chlorpheniramine	4	0	0	0	0	0	0	0	0	0	0	0	0
Citalopram	50	28	3	25	22	5	1	25	1	24	21	3	1
Clonazepam	3	2	0	2	1	1	0	2	0	2	1	1	0
Clozapine	1	1	1	0	1	0	0	1	1	0	1	0	0
Cocaine ⁴	97	64	13	51	63	0	1	64	13	51	63	0	1
Codeine ⁵	104	72	14	58	70	1	1	2	0	2	2	0	0
Cyanide	2	2	2	0	0	2	0	2	2	0	0	2	0
Cyclobenzaprine	9	6	0	6	6	0	0	4	0	4	4	0	0
Desipramine	1	1	1	0	0	1	0	1	1	0	0	1	0
Dextromethorphan	2	1	0	1	0	1	0	1	0	1	0	1	0
Diazepam	47	23	1	22	22	0	1	22	0	22	22	0	0
Difluoroethane	5	3	3	0	3	0	0	3	3	0	3	0	0
Diltiazem	1	1	0	1	1	0	0	1	0	1	1	0	0

Table 9-3 2013 Drug & Poison Caused Deaths, page 2

Drug Name	Total deaths out of 2,119 cases in which drug was present	Overdose Deaths (329) – Drug Present						Overdose Deaths (329) – Drug Causing					
		In which drug was present	Single drug OD in which drug was present	Multiple drug OD in which drug was present	Accident	Suicide	Undetermined	In which drug caused death	OD in which a single drug caused death	OD in which multiple drugs caused death	Accident	Suicide	Undetermined
Diphenhydramine	45	24	7	17	14	8	2	21	5	16	11	8	2
Doxepin	7	4	0	4	2	1	1	3	0	3	1	1	1
Doxylamine	3	2	0	2	0	2	0	2	0	2	0	2	0
Ethanol	281	105	28	77	87	17	1	81	14	67	70	10	1
Ethyl Chloride	1	1	0	1	1	0	0	1	0	1	1	0	0
Ethylene Glycol	2	2	2	0	0	2	0	2	2	0	0	2	0
Fentanyl	10	5	4	1	5	0	0	5	4	1	5	0	0
Flecainide	1	1	0	1	0	1	0	1	0	1	0	1	0
Fluoxetine	17	4	0	4	4	0	0	3	0	3	3	0	0
Gamma hydroxybutyrate	1	1	1	0	1	0	0	1	1	0	1	0	0
Hydrocodone	39	17	0	17	10	6	1	17	0	17	10	6	1
Hydromorphone	21	10	0	10	7	3	0	8	0	8	6	2	0
Imipramine	1	1	1	0	0	1	0	1	1	0	0	1	0
Isopropanol	9	2	2	0	2	0	0	0	0	0	0	0	0
Lacosamide	1	1	1	0	0	1	0	1	1	0	0	1	0
Lamotrigine	6	2	0	2	1	1	0	2	0	2	1	1	0
Lithium	1	1	0	1	0	0	1	1	0	1	0	0	1
Lorazepam	18	5	1	4	4	1	0	4	0	4	4	0	0
Meprobamate	6	4	0	4	0	0	0	0	0	0	0	0	0
Methadone	78	55	8	47	52	2	1	55	8	47	52	2	1
Methamphetamine	79	42	13	29	40	0	2	42	13	29	40	0	2
Metroprolol	2	1	0	1	0	1	0	1	0	1	0	1	0
Midazolam	34	4	2	2	4	0	0	1	0	1	1	0	0
Mirtazapine	5	3	0	3	3	0	0	2	0	2	2	0	0
Monoacetylmorphine ^b	54	52	12	40	52	0	0	0	0	0	0	0	0

Table 9-3 2013 Drug & Poison Caused Deaths, page 3

Drug Name	Total deaths out of 2,119 cases in which drug was present	Overdose Deaths (329) – Drug Present						Overdose Deaths (329) – Drug Causing					
		In which drug was present	Single drug OD in which drug was present	Multiple drug OD in which drug was present	Accident	Suicide	Undetermined	In which drug caused death	OD in which a single drug caused death	OD in which multiple drugs caused death	Accident	Suicide	Undetermined
Nortriptyline ⁷	24	15	1	14	9	6	0	2	0	2	2	0	0
Opiate ⁸	120	123	21	102	119	4	0	123	21	102	119	4	0
Oxazepam	24	11	0	11	11	0	0	1	0	1	1	0	0
Oxycodone	92	52	7	45	47	5	0	49	6	43	45	4	0
Oxymorphone	19	12	5	7	9	3	0	1	1	0	0	1	0
Paroxetine	1	1	0	1	1	0	0	1	0	1	1	0	0
Pentobarbital	3	2	0	2	1	1	0	2	0	2	1	1	0
Phenobarbital	6	1	0	1	1	0	0	1	0	1	1	0	0
Propofol	1	1	1	0	1	0	0	1	1	0	1	0	0
Psuedo/Ephedrine	1	1	0	1	1	0	0	1	0	1	1	0	0
Quetiapine	15	10	1	9	7	2	1	9	0	9	7	1	1
Temazepam	24	9	0	9	9	0	0	4	0	4	4	0	0
Topiramate	1	1	0	1	1	0	0	1	0	1	1	0	0
Tramadol	10	6	1	5	3	2	1	6	1	5	3	2	1
Trazodone	23	8	1	7	6	2	0	6	0	6	5	1	0
Venlafaxine	10	6	1	5	4	2	0	6	1	5	4	2	0
Zoplidem	27	15	0	15	12	3	0	15	0	15	12	3	0

Table 9-3 2013 Drug & Poison Caused Deaths, page 4

¹Table 9-3 is constructed on the basis of finding each of the listed drugs by laboratory analysis of the decedent's blood. The first column represents the total number of cases in which the specific drug was detected, regardless of cause and manner of death. The rest of the columns represent only drug overdose deaths and are divided into two parts. The part that lists "Drug Present" represents the number of cases in drug overdose deaths in which the drug was present in quantifiable amounts. The other part that lists "Drug Causing" represents the number of drug overdose deaths in which the specific drug caused or contributed to death in the opinion of the certifying Medical Examiner, i.e., the drug was included on the death certificate. In many cases, the numbers in the first part are more than those in the second part because the drug, although present, was not considered to contribute significantly to death, i.e., the drug was not listed on the death certificate even though it was detected in the decedent. In a few cases, the column that lists "In which drug caused death" is greater than the column that lists "In which drug was present," because the drug was detected but not in quantifiable levels, and the certifying Medical Examiner considered the drug to have contributed to death.

²Cannabinoids are listed if they were found at any level in blood or urine, not necessarily in quantified levels. Cannabinoids in levels typically found are not considered lethal agents and, therefore, there are no instances of single drug overdose deaths involving cannabinoids or THC. Although cannabinoids/THC were not considered contributory to death, they were detected in overdose deaths as listed.

³Carbon monoxide fatalities are listed in the first column if the level of carboxyhemoglobin was 5% or greater. The rest of the columns represent only drug overdose deaths and are divided into two parts, "Drug Present" and "Drug Causing". There were nine suicides from the inhalation of carbon monoxide but these deaths were listed as asphyxia deaths secondary to a suffocating and were not listed as drug overdoses. There were five accidental deaths where carbon monoxide was present. One was a drug overdose where carbon monoxide was not listed as contributing to death, three were related to fires and one was a traffic accident with fire. There were no undetermined overdose deaths involving carbon monoxide.

⁴Includes benzoylecgonine.

⁵Out of the 72 overdose deaths involving codeine, in 67 cases, the source of the drug was likely small quantities of codeine present in heroin used by illicit drug users. In 4 case the source of the drug was likely pharmaceutical codeine and one case was from an unknown source.

⁶ Monoacetylmorphine (MAM) is a principal toxicological marker for heroin. It is the first breakdown product of heroin, which is diacetylmorphine. The presence of MAM, therefore, proves the source of opiate to be heroin. However, the absence of MAM does not imply that the source of the opiate was not heroin.

⁷In 5 of the 15 total cases, nortriptyline was present without the presence of amitriptyline, indicating that the source of the drug was, in fact, nortriptyline. In the other 10 cases, amitriptyline was also present, indicating that the nortriptyline was present due to the breakdown of amitriptyline. There were a total of 2 nortriptyline overdose deaths; both were accidental multiple drug overdoses.

⁸As used in this section, "opiate" refers exclusively to the naturally occurring drug (morphine) or its derivative (heroin). This category does not include the other "opioids" such as oxycodone, hydrocodone, hydromorphone, oxymorphone and methadone. In 2013 there were 123 deaths caused by opiates. Toxicological analysis detects only morphine and cannot differentiate heroin and pharmaceutical morphine as the likely source of the opiate. Based on toxicology analysis (presence of acetylmorphine), scene investigation, and circumstances it was determined that out of these 123 deaths, 94 were definitely or probably due to heroin and 17 were due to pharmaceutical morphine. In the remaining 12 cases it was not possible to determine whether the death was due to heroin or pharmaceutical morphine.

Graph 9-1 Drug & Poison Caused Deaths / Accident, Suicide, Undetermined / KCME / 2003- 2013




Table 9-4 Total Overdose Deaths / Accident, Suicide, Undetermined / 2003 – 2013⁹


Overdose Deaths	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Accident	211	216	262	247	232	233	217	205	230	279
Suicide	41	39	36	36	29	29	43	48	51	41
Undetermined	26	18	14	19	17	9	11	15	17	9
Totals	278	273	312	302	278	271	271	268	298	329

⁹ Includes all deaths classified as overdose, regardless of whether lab samples were available for analysis.

Table 9-5 Overdose Deaths Caused by Cocaine, Methadone, Opiates, Methamphetamine, or Oxycodone¹⁰ / KCME / 2004 - 2013


DRUG	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Cocaine	93	90	114	89	74	62	48	44	48	64
Methadone	68	86	96	84	86	85	67	54	56	52
Methamphetamine	18	29	21	21	15	19	19	22	40	42
Opiates	89	88	83	87	83	71	72	87	103	170
Oxycodone	32	34	47	51	44	60	35	42	36	49

Graph 9-2 Drug & Poison Caused Deaths / Accident, Suicide, Undetermined / KCME / 2003 - 2013




¹⁰ In this context, "caused by" refers to single or multiple drug overdoses in which the drug was listed on the death certificate.

Graph 9-3 Cocaine Involved Deaths¹¹ / KCME / 2004 – 2013




Graph 9-4 Ethanol Involved Deaths / KCME/ 2004– 2013




¹¹In Graphs 9-3, 9-4, 9-5 and 9-6, "overdose" refers to deaths due to the listed drug or ethanol in single or multiple drug overdose deaths where the listed drug or ethanol was listed on the death certificate.


Graph 9-5 Methadone Involved Deaths / KCME / 2004 - 2013




Graph 9-6 Methamphetamine Involved Deaths / KCME / 2004 – 2013



Graph 9-7 Opiate Overdose Deaths & Heroin-Related Deaths / KCME / 2004 - 2013¹²



Graph 9-8 Oxycodone Involved Deaths / KCME / 2004- 2013



¹²In 2004, the King County Medical Examiner's Office began collecting data on probable heroin overdoses based on a combination of circumstances, scene investigation and toxicology results.

Graph 9-9 Drug / Poison Deaths / Age / KCME / 2004 – 2013






Table 9-6 Drug / Poison Deaths / Age / KCME / 2013

AGE GROUP (YEARS) / GENDER		MANNER OF DEATH			SUB-TOTAL	TOTAL
		ACCIDENT	SUICIDE	UNDETERMINED		
<20		2	1	2		5
Male		2	1	1	4	
Female		0	0	1	1	
20-29		41	2	0		43
Male		28	1	0	29	
Female		13	1	0	14	
30-39		58	6	4		68
Male		45	1	3	49	
Female		13	5	1	19	
40-49		69	13	3		85
Male		47	6	2	55	
Female		22	7	1	30	
50-59		76	8	0		84
Male		42	3	0	45	
Female		34	5	0	39	
60-69		29	9	0		38
Male		19	5	0	24	
Female		10	4	0	14	
70-79		4	1	0		5
Male		4	0	0	4	
Female		0	1	0	1	
80-89		0	0	0		0
Male		0	0	0	0	
Female		0	0	0	0	
90+		0	1	0		1
Male		0	1	0	1	
Female		0	0	0	0	
Totals		279	41	9		329

Deaths due to firearms

The Medical Examiner is responsible for investigating all deaths due to firearms that occur in King County. Medical Examiner data relate primarily to the victim because information regarding the weapon and the shooter is often unknown. The following data are specific to the victims of firearm deaths.

In 2013, the Medical Examiner investigated 145 firearm deaths. In 2012, firearms caused 169 deaths. Of the 145 firearm deaths in 2013, 44 (30%) were homicides and 100 (69%) were suicides. One firearm death was classified as an accident in 2013. In 2012, there were two firearm deaths classified as accidents. In 2013, there were no firearm deaths that were classified as undetermined; there were two in 2012.

In 2013, gunshot wounds were the leading cause of death for homicides and suicides. Firearm deaths comprised 59% (44/74) of homicides, compared to 68% (47/69) in 2012. In 2013, suicides by firearms represented 38% (100/266) of suicide deaths compared to 42% (119/281) in 2012.

In 2013, of the 44 firearm homicide victims, 9% (4/45) were 19 years old and younger – a decrease from 2012 when 15% of firearm homicide victims were 19 years old and younger. In 2013, it is estimated that a disproportionate number of firearm homicide victims were African American (47%, 21/45) compared to the percentage of African Americans in the general population (see discussions on pages 8 and 44). Of the 21 African American firearm homicide victims, three were males 19 years old and younger and nine were males between 20 and 29 years of age. In comparison, 38% (17/45) of the homicide firearm victims were white. Of the 17 white homicide victims, 53% (9/17) were males between 20 and 29 years old.

Of the 100 firearm suicide victims in 2013, 85% (85/100) were white and 74% (74/100) were males. One of the firearm suicide victims was 19 years old or under (1%, 1/100). Thirty (30%, 30/100) of the gunshot suicide victims were between the ages of 20 and 39 years of age, 26 (26%, 26/100) were between 40 and 59 years, and 43 (43%, 43/100) were 60 years and older.



Table 10-1 Firearm Deaths / Manner / Age / Gender / KCME / 2013

AGE GROUP / GENDER	MANNER OF DEATH				SUB-TOTAL	TOTAL
	A	H	S	U		
<13 years	0	0	0	0		0
<i>Male</i>	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	
13-15 years	0	0	0	0		0
<i>Male</i>	0	0	0	0	0	
<i>Female</i>	0	0	0	0	0	
16-19 years	0	4	1	0		5
<i>Male</i>	0	4	1	0	5	
<i>Female</i>	0	0	0	0	0	
20-29 years	0	22	15	0		37
<i>Male</i>	0	19	12	0	31	
<i>Female</i>	0	3	3	0	6	
30-39 years	0	10	15	0		25
<i>Male</i>	0	10	13	0	23	
<i>Female</i>	0	0	2	0	2	
40-49 years	1	3	7	0		11
<i>Male</i>	1	3	6	0	10	
<i>Female</i>	0	0	1	0	1	
50-59 years	0	4	19	0		23
<i>Male</i>	0	4	16	0	20	
<i>Female</i>	0	0	3	0	3	
60-69 years	0	1	23	0		24
<i>Male</i>	0	1	19	0	20	
<i>Female</i>	0	0	4	0	4	
70-79 years	0	0	9	0		9
<i>Male</i>	0	0	9	0	9	
<i>Female</i>	0	0	0	0	0	
80-89 years	0	0	8	0		8
<i>Male</i>	0	0	8	0	8	
<i>Female</i>	0	0	0	0	0	
90+	0	0	3	0		3
<i>Male</i>	0	0	3	0	3	
<i>Female</i>	0	0	0	0	0	
Totals	1	44	100	0		145
Percent	1%	30%	69%	0%		100%

Graph 10-1 Firearm Deaths / Manner / Age Group / KCME / 2013






Table 10-2 Firearm Deaths / Manner / Race / Gender / KCME / 2013

RACE / GENDER	MANNER OF DEATH				SUB-TOTAL	TOTAL
	A	H	S	U		
Asian/Pacific Islander	0	7	4	0		11
<i>Male</i>	0	7	4	0	11	
<i>Female</i>	0	0	0	0	0	
African American	0	21	6	0		27
<i>Male</i>	0	20	4	0	24	
<i>Female</i>	0	1	2	0	3	
Am Indian / AK Native	0	2	1	0		3
<i>Male</i>	0	2	1	0	3	
<i>Female</i>	0	0	0	0	0	
White	1	17	85	0		103
<i>Male</i>	1	16	74	0	91	
<i>Female</i>	0	1	11	0	12	
Other	0	1	0	0		1
<i>Male</i>	0	0	0	0	0	
<i>Female</i>	0	1	0	0	1	
Totals	1	44	100	0		145
Percents	1%	30%	69%	0%		100%

Causes of death in children and youth

In 2013, the King County Medical Examiner's Office investigated 83 deaths of children and youth ages 19 years or younger, which represented 4% (83/2,119) of the total deaths investigated. Of these deaths, 27% (22/83) were natural, 13% (11/83) were accidental (non-traffic), 10% (8/83) were homicides, 16% (13/83) were traffic-related, 18% (15/83) were suicides, and 17% (14/83) were classified as manner undetermined. In addition to investigating childhood deaths, the King County Medical Examiner participates in Child Death Review, a process which discusses these deaths in detail and formulates prevention strategies.

Of the 22 natural deaths of children and youth investigated by the Medical Examiner, 59% (13/22) were of infants less than one year of age. Of these 13 infants who died of natural causes, 7 were due to Sudden Infant Death Syndrome (SIDS). In addition, 10 infant deaths were classified as "Sudden Unexplained Infant Death" (SUID), manner undetermined, due to the inability to exclude if external factors contributed to death.

There were 13 homicides among children and youth. Of these 13 homicide victims, 10 were teenagers (13 - 19 years of age), 1 was a child (one to 12 years of age), and 1 was an infant less than one year of age. Homicides as a result of gunshot wounds accounted for 31% (4/13) of the children and youth homicide victims.

There were 15 youth suicides, with the majority, 13, being between the ages of 12 and 19 years. 2 suicides occurred in the age group of 1 – 12 years of age. Males comprised 87% (13/15) of the victims. Regarding the methods used to commit suicide by youth, 1 was by firearm, 11 were by hanging, 1 was drugs / alcohol and 1 was from inhalation of carbon monoxide, and one was from a jump.

13 children and youth (19 years and under) died in traffic-related accidents, of whom 12/13 (92%) were teenagers 13 – 19 years of age. There were 7 motor vehicle driver deaths, 3 motor vehicle passenger deaths, 1 pedestrian death, 1 motorcycle driver death, and 1 skateboarder in traffic death. Of the 10 children and youth who died in automobiles, 4 were known to be restrained, 4 unrestrained and 2 were restraint unknown.

The following tables list the causes of death among children and youth for all manners in three age groups: less than one year, 1-12 years and 13-19 years.

Graph 11-1 Causes of Death in Children & Youth / KCME / 2013




Table 11-1 Causes of Death: Children Under 1 Year of Age / KCME / 2013

CIRCUMSTANCES	MANNER OF DEATH						SUB-TOTAL	TOTAL
	A	H	S	T	U	N		
Miscellaneous								22
Asphyxia	1	0	0	0	0	0	1	
Drowning	0	0	0	0	0	0	0	
Prematurity	0	0	0	0	0	0	0	
Other	0	2	0	0	0	0	2	
SIDS	0	0	0	0	12 ²²	7	19	
Other Natural Disease	0	0	0	0	0	6		6
Totals	1	2	0	0	12	13		28

²² Includes 10 cases classified as Sudden Unexplained Infant Death with the possibility of bed sharing listed as a significant condition contributing to the death.

Table 11-2 Causes of Death: Children 1 to 12 Years of Age / KCME / 2013

CIRCUMSTANCES	MANNER OF DEATH						SUB-TOTAL	TOTAL
	A	H	S	T	U	N		
Asphyxia	2	0	2	0	0	0		4
<i>Carbon Monoxide</i>	0	0	0	0	0	0	0	
<i>Drowning</i>	1	0	0	0	0	0	1	
<i>Hanging</i>	0	0	2	0	0	0	2	
<i>Mechanical</i>	1	0	0	0	0	0	1	
<i>Other</i>	0	0	0	0	0	0	0	
<i>Positional</i>	0	0	0	0	0	0	0	
Miscellaneous	2	0	0	0	0	0		2
<i>Complication of Therapy</i>	0	0	0	0	0	0	0	
<i>Drugs / Poisons</i>	0	0	0	0	0	0	0	
<i>Fall</i>	1	0	0	0	0	0	1	
<i>Fire / Explosion</i>	1	0	0	0	0	0	1	
<i>Hyperthermia</i>	0	0	0	0	0	0	0	
<i>Jump</i>	0	0	0	0	0	0	0	
<i>Non Traffic -Vehicle</i>	0	0	0	0	0	0	0	
<i>Other</i>	0	0	0	0	0	0	0	
Physical Trauma	0	1	0	0	1	0		2
<i>Abuse</i>	0	0	0	0	0	0	0	
<i>Blunt Force / Crushing</i>	0	1	0	0	1	0	2	
<i>Burns / Fire</i>	0	0	0	0	0	0	0	
<i>Firearms</i>	0	0	0	0	0	0	0	
<i>Incised / Stab Wound(s)</i>	0	0	0	0	0	0	0	
<i>Other</i>	0	0	0	0	0	0	0	
Transportation Related	0	0	0	1	0	0		1
<i>Bicycle</i>	0	0	0	0	0	0	0	
<i>Motor Vehicle Driver</i>	0	0	0	0	0	0	0	
<i>Motor Vehicle Passenger</i>	0	0	0	1	0	0	1	
<i>Other</i>	0	0	0	0	0	0	0	
<i>Pedestrian</i>	0	0	0	0	0	0	0	
Natural Disease	0	0	0	0	0	6		6
Totals	4	1	2	1	1	6		15

Table 11-3 Causes of Death: Children 13 to 19 Years of Age / KCME / 2013

CIRCUMSTANCES	MANNER OF DEATH						SUB-TOTAL	TOTAL
	A	H	S	T	U	N		
Asphyxia	3	0	10	0	0	0		13
<i>Carbon Monoxide</i>	0	0	1	0	0	0	1	
<i>Drowning</i>	3	0	0	0	0	0	3	
<i>Hanging</i>	0	0	9	0	0	0	9	
<i>Smothering</i>	0	0	0	0	0	0	0	
<i>Positional</i>	0	0	0	0	0	0	0	
<i>Other</i>	0	0	0	0	0	0	0	
Drugs / Alcohol	2	0	1	0	0	0		3
Miscellaneous	0	0	1	0	0	0		1
<i>Complication of Therapy</i>	0	0	0	0	0	0	0	
<i>Fall</i>	0	0	0	0	0	0	0	
<i>Jump</i>	0	0	1	0	0	0	1	
<i>Non-Traffic Vehicular</i>	0	0	0	0	0	0	0	
<i>Other</i>	0	0	0	0	0	0	0	
Physical Trauma	1	5	1	0	1	0		8
<i>Blunt Force / Crushing</i>	1	1	0	0	1	0	3	
<i>Burns / Fire</i>	0	0	0	0	0	0	0	
<i>Firearms</i>	0	4	1	0	0	0	5	
<i>Homicidal Violence</i>	0	0	0	0	0	0	0	
<i>Incised / Stab Wound(s)</i>	0	0	0	0	0	0	0	
<i>Strangulation</i>	0	0	0	0	0	0	0	
Transportation Related	0	0	0	12	0	0		12
<i>Bicycle</i>	0	0	0	0	0	0	0	
<i>Motor Vehicle Driver</i>	0	0	0	7	0	0	7	
<i>Motor Vehicle Passenger</i>	0	0	0	2	0	0	2	
<i>Motorcycle</i>	0	0	0	1	0	0	1	
<i>Pedestrian</i>	0	0	0	1	0	0	1	
<i>Other</i>	0	0	0	1	0	0	1	
Natural Disease	0	0	0	0	0	3		3
Totals	6	5	13	12	1	3		40

Organ donation

Although the King County Medical Examiner's Office does not approach families for donation of organs and tissue from decedents, we recognize the tremendous need for this life-saving activity and cooperate fully with organ and tissue procurement agencies. It is the philosophy of the King County Medical Examiner's Office that all requests for organ and/or tissue donation be given high priority for approval. In practice, the procurement agency contacts the KCMEO with information regarding a potential donor and the specific organs or tissue requested. The Medical Examiner then evaluates the request to determine if the donation would significantly affect the postmortem examination. In the great majority of cases, examinations can be conducted so that donations do not interfere with certification of death or collection of evidence. In this way, the King County Medical Examiner's Office works to maximize the donation of organs and tissue that go directly to save lives.

In 2013, the King County Medical Examiner's Office gave release on 32 deaths that came under the office's jurisdiction. Altogether, there were 111 organs donated for transplant from the 32 cases referred to the King County Medical Examiner. The number of specific organs transplanted in 2013 is shown in Table 12-1. In addition to the living organs listed in Table 12-1 that were donated in 2013, the KCMEO approved the donation of skin, bone, cartilage, heart valves, corneas and other tissues through the tissue procurement agency, Northwest Tissue Service. Altogether, there were 74 donors who, on average, were able to provide over 50 donations each (3,700 total) to tissue transplant recipients.

ORGAN	# Transplanted
Heart	12
Intestine	0
Kidney	60
Liver	19
Lung	15
Pancreas	5
Total	111

Disposition review

All deaths covered under RCW 68.50.010 are required by law to be reported to the Medical Examiner, however in the past these deaths have not always been reported in a timely manner. For some of these deaths, a complete investigation is not possible because the body was cremated prior to the death being reported to the Medical Examiner.

Beginning January 1, 2008, the King County Council authorized the Medical Examiner's Office to review the death certificates of all decedents to be cremated in order to rule out the need for additional investigation and ensure the proper determination of cause and manner of death.

In 2013, the Medical Examiner's Office handled 10,597 cremation review requests. In 67 cases the Medical Examiner took jurisdiction to investigate further and determine correct cause and manner of death. Without this cremation review, these cases would not have been seen and the correct determination of death missed.

Beginning January 1, 2013, the King County Council authorized the Medical Examiner's Office to review the death certificates of all decedents to be buried in order to rule out the need for additional investigation and ensure the proper determination of cause and manner of death.

In 2013, the Medical Examiner's Office handled 3,329 burial review requests. In 11 cases the Medical Examiner took jurisdiction to investigate further and determine correct cause and manner of death. Without this burial review, these cases would not have been seen and the correct determination of death missed.

Medical Examiner activity

The staff of the Medical Examiner's Office are involved in a wide variety of activities commensurate with the mission of the office including responding to and investigating the scene of death, performing postmortem examinations, certifying the cause and manner of death, and providing information and assistance to families. Investigators, who are familiar with the emotional trauma of an unexpected death, communicate directly with families as do the Medical Examiner pathologists, who review their findings with the families in order to clarify the many questions that accompany a sudden loss of life. The office also provides referrals to grief support services.

In all cases investigated by the Medical Examiner, it is essential that the decedent's identity is established and the next-of-kin is located and notified regarding the death. In addition, property belonging to the decedent must be controlled and released according to legal requirements. In most cases these issues are resolved expeditiously. In certain cases, identification requires additional effort in locating dental, medical or police records. Some individuals may have died leaving no next-of-kin or next-of-kin far removed. Ensuring that all leads have been exhausted in pursuit of next-of-kin can be a very time consuming but ultimately a rewarding effort.

The postmortem examination on each decedent includes the preservation of various body fluids and tissues for microscopic and toxicologic analysis. Photographs are taken of the external and internal portions of the examination, which are available for review at a later date if needed. Photographic documentation is also an essential item in those cases where the pathologist must provide court testimony. Forensic Anthropology is another important activity necessary to resolve skeletal cases and difficult identification issues.

Medical Examiner pathologists, anthropologist and investigators provide testimony in court and at depositions. Staff participates in meetings with police, medical professionals, and attorneys. A recent addition to the duties of the Chief Medical Examiner is expert medical consultation and testimony in cases involving nonfatal domestic violence assaults.

Autopsy reports and related data from individual investigations are provided to law enforcement agencies, prosecuting attorneys and many other agencies including Labor and Industries, the Drug Enforcement Administration, and the Consumer Product Safety Commission. Drug deaths are reported to the Drug Abuse Warning Network (DAWN).

In 2003, the Medical Examiner's Office created a student internship program that provides educational opportunities for students interested in forensic autopsy and death investigation. Through this program, numerous interns have obtained full-time careers in death investigation, both at the KCMEO and in other area medical examiner's offices.

Medical Examiner investigations require frequent contact between the Medical Examiner's Office and the news media. Staff members are skilled in responding to the media inquiries that occur daily. The Medical Examiner pathologists and other staff participate in a variety of medical conferences, and provide information on a regular basis to law enforcement and to medical personnel on various aspects regarding the role and function of the Medical Examiner's Office.

The data collected and presented in this and other Medical Examiner annual reports also provide baseline information for further analysis. Medical Examiner staff analyzes data to study relevant death investigation topics that have applications in such fields as law enforcement, medicine, law, social sciences, and injury prevention. Examples include infant mortality, teenage suicide, child abuse, law enforcement restraint, investigation of vehicular traffic collisions, and investigation of therapeutic complication deaths. In addition, the office participates in teaching medical students, pathology residents, emergency medical service, and law enforcement personnel.

In 2013, staff participated as speakers at universities, conferences, and training seminars for law enforcement, medical, legal, and social service personnel in the following presentations and lectures:

Richard C. Harruff, M.D., Ph.D., Chief Medical Examiner

Academic appointment

- Clinical Associate Professor, Department of Pathology, University of Washington School of Medicine.

Professional organizations

- American Academy of Forensic Sciences.
- National Association of Medical Examiners.
- Disaster Mortuary Operations Response Team, Region 10.

Preceptor and faculty positions

- Program Director, King County Medical Examiner's Office Fellowship Training Program in Forensic Pathology.
- University of Washington School of Medicine medical students and pathology residents.
- Course Director and Faculty, "Problems in Forensic Pathology", King County Medical Examiner's Office.
- Host for one-year International Scholar from Sri Lanka

Scientific publication

- Harruff RC, Park J, Smelser BJ. Relation of kinetic energy to contact wounds of the head by center fire rifles and shotgun slugs. J Forensic Sci. 2013 Jan;58(1):69-72.

Educational presentations

- Introduction to the King County Medical Examiner's Office. University of Washington and Harborview Medical Center Paramedic Training, Seattle, Washington, March 14.

- Strangulation and pattern injuries. Harborview Center for Sexual Assault and Traumatic Stress, Sexual Assault Nurse Examiner Training, Seattle, Washington, March 20.
- Medicolegal investigation of deaths in infants and young children. Training for Office of the Public Defender, Seattle, Washington, March 29.
- Medicolegal death investigation. Presentation for The Defender Association, Seattle, Washington, April 30.
- Investigation of traffic fatalities. Training for traffic investigator's course, Seattle Police Department, Seattle, Washington, May 1.
- Medicolegal death investigation. University of Washington Private Investigators Course, Seattle, Washington, June 7.
- Mass fatality management: scene recovery. 2013 Integrated Training – Region 10 Disaster Mortuary Operational Response Team, National Disaster Medical System, and Fatality Search and Recovery Team, Washington Air National Guard, Camp Murray, Washington, September 7.
- Mechanism of injury in traffic fatalities. Traffic Collision Reconstruction Course, Washington State Patrol, Shelton, Washington, October 17.
- Medicolegal Investigation of Deaths in the Vulnerable and Elderly. Presentation for Washington State's 10th Annual Conference on Abuse of Elders and Adults with Disabilities: "Improving Our Response to Neglect and Self-Neglect", Burien, Washington, October 29.
- Contemporary US education in forensic pathology: core competencies and milestones. College of Forensic Pathologists of Sri Lanka, Colombo, Sri Lanka, December 10
- Medicolegal investigation of deaths in elderly and vulnerable adults. Medicolegal Society of Sri Lanka, Colombo, Sri Lanka, December 12

Miscellaneous activities

- Medical examiner responsibilities in deaths involving Somali Muslims. Discussion with Somali Health Board, Seattle, Washington, April 24.
- External examiner for MD (Forensic Medicine) Examinations, Colombo, Sri Lanka, December 10-14

Aldo Fusaro, DO, Associate Medical Examiner

Academic Appointment

- Clinical Assistant Professor, Department of Pathology, University of Washington School of Medicine

Preceptorship

- University of Washington School of Medicine, medical students and pathology residents
- King County Medical Examiner's Office, forensic pathology fellow trainer

Associations, Committees and Boards

- Member, American Medical Association
- Member, Washington Association of Coroners and Medical Examiners

- Member, Washington State Medical Association
- Member, National Association of Medical Examiners
 - Membership Committee
 - Delegate to the American Medical Association
- Fellow, College of American Pathologists
- Fellow, American College of Clinical Pathologists
- Advisory Committee, King County Medical Examiner's Office
- Child Death Review Committee, King County Medical Examiner's Office
- Elder Death Review Committee, King County Medical Examiner's Office
- Quality Improvement Subcommittee, King County Medical Examiner's Office
- Multiple Fatality Incident Preparedness Team, Seattle King County Public Health

Professional Meetings, Trainings and Certifications

- Forensic Investigations Council Meetings- January, March, April, September
- Annual Blood Borne Pathogens Training, Public Health- Seattle and King County, September
- Health Information Privacy and Security Training, Public Health- Seattle and King County, December

Local and Regional Educational Presentations:

- Cases in Forensic Pathology (2),
Eastside Preparatory School - Kirkland, WA - April-May, 2013
- What is a SUID?
Regional Justice Training Center - Burien, WA - May, 2013
- Basic Homicide Investigation - Attorney General's Office
Regional Justice Training Center - Burien, WA - June, 2013
- Forensic Case Studies
Sammamish High School - Bellevue, WA - September, 2013

Katherine M. Taylor, Ph.D., D-ABFA Forensic Anthropologist

Academic Affiliation

- University of Washington Department of Anthropology: Affiliate Faculty
- Seattle University Department of Criminal Justice: Adjunct Faculty

Associations, Committees, and Boards

- Fellow, American Academy of Forensic Sciences
- Diplomat, American Board of Forensic Anthropology
- Member, Society of Forensic Anthropologists
- Member, Seattle University Criminal Justice Advisory Board
- Member, Bellevue College Criminal Justice Advisory Board

Educational Presentations

- “Forensic Anthropology in Homicide Investigations.” A presentation to the Basic Homicide Investigation class sponsored by the Washington State Attorney General’s Office. Burien WA, June 11th.

Greg Hewett, Mdiv, Administrator

Associations, Committees & Boards

- Member, Seattle University Advisory Committee, Criminal Justice Program
- Member, Washington Association of Coroners and Medical Examiners

Educational Presentation

Group Health staff training. Nurses reporting requirements. Seattle, Washington, October 9th

David Delgado, D-ABMDE, Medicolegal Investigator II

Association

- Diplomate, American Board of Medicolegal Death Investigators
- Member, Washington Associations of Coroners and Medical Examiners

Educational Presentation

Sudden Unexpected Infant Death and Child Fatality Investigations, Washington State Criminal Justice Training Commission, Burien, WA; May 2013

William Barbour, BS, D-ABMDI, Medicolegal Investigator II

Associations

- Diplomate, American Board of Medicolegal Death Investigators
- Member, Washington Associations of Coroners & Medical Examiners

Educational Presentations

- Role and Responsibility of the King County Medical Examiner's Office
- Seattle University Biology - KCMEO - Seattle, WA January 24
- Seattle University Biology - KCMEO - Seattle, WA January 31
- Seattle University Biology – KCMEO- Seattle, WA February 7
- Shoreline Police Department Volunteers – KCMEO – Seattle, WA August 28
- Seattle University Biology – KCMEO – Seattle, WA November 6
- Seattle University Criminal Justice Club – Seattle, WA November 20

Jonathan Z. Gallar, BCJ, D-ABMDI, Medicolegal Investigator I

Educational Presentations

- Medicolegal Miscellanea – KCMEO – Seattle, WA October 23

Kyle Schwab, MCJ, D-ABMDI, Medicolegal Investigator I

Educational Presentations

- Role and Responsibility of the King County Medical Examiner's Office
- Seattle University Biology - KCMEO - Seattle, WA January 24
- Seattle University Biology - KCMEO - Seattle, WA January 31
- Seattle University Biology – KCMEO- Seattle, WA February 7
- Shoreline Police Department Volunteers – KCMEO – Seattle, WA August 28
- Seattle University Biology – KCMEO – Seattle, WA November 6
- Seattle University Criminal Justice Club – Seattle, WA November 20

Presentations/Guest Speaking

- Seattle University: CRJS Forensic Science – Jan 24
- Seattle University: KCME Presentation/Tour – May 1
- KCME Wednesday Conference – Adolescent suicide – July 17
- DMORT X training – death investigations during mass fatality – Fort Lewis – September 7
- KCME Wednesday Conference – Medicolegal Miscellanea: Knives, Axes, and Saws – October 23

Associations

- Seattle University Criminal Justice Advisory Committee
- ABMDI
- King County Public Health Reserve Corps

Written work

- Guns and Death: A frequency analysis of firearm related deaths within King County, WA in 2011 (Jan, 2013)

Samantha Jasso, B.J.C., Health Program Assistant I

Educational Presentations

- RCPGP Family Assistance Center Seminar and Tabletop Exercise – Recap, Seattle, WA – March 20
- Mass Fatality VIP Program: An Interactive Primer – Seattle, WA – April 24
- Recovery Scenarios – Fort Lewis, WA – September 7
- Exploring Homicide-Suicides: A Closer Look at Incidents in King County – KCMEO – Seattle, WA – October 16




Table 14-1 Weekly Variation of Deaths Investigated by the King County Medical Examiner's Office

	TOTAL
Number of weeks studied	52
Mean number of ME jurisdiction cases per week	41
Maximum ME jurisdiction cases in any one week	53
Minimum ME jurisdiction cases in any one week	25

Table 14-2 Weekly Variation of Autopsies Investigated by the King County Medical Examiner's Office

	TOTAL
Number of weeks studied	52
Mean number of autopsies performed per week	26
Maximum # autopsies performed in any one week	40
Minimum # autopsies performed in any one week	14

Organization of the King County Medical Examiner's Office 2013



Glossary of Terms

Blood alcohol level:

The concentration of ethanol (alcohol) found in blood following ingestion. Measured in grams per 100 ml of blood or grams %. In the State of Washington, 0.08 grams % is considered the legally intoxicated level while driving.

Cause of Death:

Any injury or disease that produces a physiological derangement in the body that results in the death of an individual.¹

Drug:

Therapeutic drug: A substance, other than food, used in the prevention, diagnosis, alleviation, treatment, or cure of disease.

Recreational drug: A drug used non-medically for personal stimulation/depression/euphoria.

Drug-caused death:

Death directly caused by a drug or drugs in combination with each other or with alcohol.

Fetal Death:

Category of deaths that occur within the uterus. The Medical Examiner assumes jurisdiction over fetal deaths that meet the criteria specified in RCW 68.50. See pages 2 - 3 of this report for details.

Jurisdiction:

The jurisdiction of the Medical Examiner extends to all reportable deaths occurring within the boundaries of King County, whether or not the incident leading to the death (such as an accident) occurred within the county. Reportable deaths are defined by RCW 68.50, as explained in the "Description and Purpose" section of this report. Not all natural deaths reported fall within the jurisdiction of the Medical Examiner.

Manner of Death:

A classification of the way in which the events preceding death were causal factors in the death. The manner of death as determined by the forensic pathologist is an opinion based on the known facts concerning the circumstances leading up to and surrounding the death, in conjunction with autopsy findings and laboratory tests.²

¹DiMaio, Vincent J. & DiMaio, Dominick. Forensic Pathology, Second Edition. CRC Press, 2001.

²Ibid, p. 3

Manner: Accident

Death other than natural, where there is no evidence of intent, i.e., unintentional. In this report, traffic accidents are classified separately.

Manner: Homicide

Death resulting from intentional harm (explicit or implicit) of one person by another, including actions of grossly reckless behavior.

Manner: Natural

Death caused solely by disease. If natural death is hastened by injury (such as a fall or drowning in a bathtub), the manner of death is classified other than natural. The Natural category includes complication of therapy deaths.

Manner: Suicide

Death as a result of a purposeful action with intent (explicit or implicit) to end one's own life.

Manner: Traffic

Unintentional deaths of drivers, passengers, and pedestrians involving motor vehicles on public roadways. Accidents involving motor vehicles on private property (such as driveways) are not included in this category and are classified non-traffic, vehicular accidents.

Manner: Undetermined

Manner assigned when there is insufficient evidence or information, especially about intent, to assign a specific manner.

Opiate:

Any preparation or derivative of opium, including heroin, morphine or codeine. In this report "opiate deaths" most likely refer to heroin caused deaths.

Poison:

Any substance, either taken internally or applied externally, that is injurious to health or dangerous to life, and with no medicinal benefit.



Race:

The racial categories used in this report are: White, African American, American Indian/Alaska Native, Asian/Pacific Islander, and Other.